6 □ APPLICATIONS OF INTEGRATION

6.1 Areas Between Curves

1. (a) $A = \int_{x=0}^{x=2} (y_T - y_B)\,dx = \int_0^2 [(3x - x^2) - x]\,dx = \int_0^2 (2x - x^2)\,dx$

(b) $\int_0^2 (2x - x^2)\,dx = \left[x^2 - \frac{1}{3}x^3\right]_0^2 = \left(4 - \frac{8}{3}\right) - 0 = \frac{4}{3}$

2. (a) $A = \int_{x=0}^{x=1} (y_T - y_B)\,dx = \int_0^1 (e^x - x^2)\,dx$

(b) $\int_0^1 (e^x - x^2)\,dx = \left[e^x - \frac{1}{3}x^3\right]_0^1 = \left(e - \frac{1}{3}\right) - (1 - 0) = e - \frac{4}{3}$

3. (a) $A = \int_{y=-1}^{y=1} (x_R - x_L)\,dy = \int_{-1}^1 [e^y - (y^2 - 2)]\,dy = \int_{-1}^1 (e^y - y^2 + 2)\,dy$

(b) $\int_{-1}^1 (e^y - y^2 + 2)\,dy = \left[e^y - \frac{1}{3}y^3 + 2y\right]_{-1}^1 = \left(e^1 - \frac{1}{3} + 2\right) - \left(e^{-1} + \frac{1}{3} - 2\right) = e - \frac{1}{e} + \frac{10}{3}$

4. (a) $A = \int_{y=0}^{y=3} (x_R - x_L)\,dy - \int_0^3 [(2y - y^2) - (y^2 - 4y)]\,dy = \int_0^3 (-2y^2 + 6y)\,dy$

(b) $\int_0^3 (-2y^2 + 6y)\,dy = \left[-\frac{2}{3}y^3 + 3y^2\right]_0^3 = (-18 + 27) - 0 = 9$

5. $A = \int_{-2}^0 [(x^3 - 3x) - x]\,dx + \int_0^2 [x - (x^3 - 3x)]\,dx \overset{\text{symmetry}}{=} 2\int_0^2 [x - (x^3 - 3x)]\,dx$

$= 2\int_0^2 (4x - x^3)\,dx = 2\left[2x^2 - \frac{1}{4}x^4\right]_0^2 = 2[(8 - 4) - 0] = 8$

6. $A = \int_{-2}^1 \left[\left(\frac{2}{3}x + \frac{16}{3}\right) - x^2\right]dx + \int_1^2 [(-2x + 8) - x^2]\,dx$

$= \left[\frac{1}{3}x^2 + \frac{16}{3}x - \frac{1}{3}x^3\right]_{-2}^1 + \left[-x^2 + 8x - \frac{1}{3}x^3\right]_1^2$

$= \left[\left(\frac{1}{3} + \frac{16}{3} - \frac{1}{3}\right) - \left(\frac{4}{3} - \frac{32}{3} + \frac{8}{3}\right)\right] + \left[\left(-4 + 16 - \frac{8}{3}\right) - \left(-1 + 8 - \frac{1}{3}\right)\right] = \frac{44}{3}$

7. By inspection, we see that the curves intersect at $x = 0$.

$A = \int_0^1 (3^x - 2^x)\,dx$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. By inspection, we see that the curves intersect at $x = 1$.

$$A = \int_1^2 [\ln(x^2) - \ln x]\,dx$$



9. The curves intersect when $2 - x = 2x - x^2 \quad \Leftrightarrow$

$x^2 - 3x + 2 = 0 \quad \Leftrightarrow \quad (x-2)(x-1) = 0 \quad \Leftrightarrow$

$x = 1$ or $x = 2$.

$$A = \int_1^2 [(2x - x^2) - (2 - x)]\,dx$$



10. The curves intersect when $y^4 = 2 - y^2 \quad \Leftrightarrow$

$y^4 + y^2 - 2 = 0 \quad \Leftrightarrow \quad (y^2+2)(y^2-1) = 0 \quad \Leftrightarrow$

$y = \pm 1$.

$$A = \int_{-1}^1 [(2 - y^2) - y^4]\,dy$$



11. $A = \displaystyle\int_0^1 [(x^2+2) - (-x-1)]\,dx$

$= \displaystyle\int_0^1 (x^2 + x + 3)\,dx$

$= \left[\frac{1}{3}x^3 + \frac{1}{2}x^2 + 3x\right]_0^1 = \left(\frac{1}{3} + \frac{1}{2} + 3\right) - 0$

$= \dfrac{23}{6}$



12. $A = \displaystyle\int_{-1}^0 [(2-x) - (1+x^3)]\,dx$

$= \displaystyle\int_{-1}^0 (1 - x - x^3)\,dx$

$= \left[x - \frac{1}{2}x^2 - \frac{1}{4}x^4\right]_{-1}^0$

$= \left[0 - \left(-1 - \frac{1}{2} - \frac{1}{4}\right)\right]$

$= \dfrac{7}{4}$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. $A = \int_1^2 \left(\frac{1}{x} - \frac{1}{x^2}\right) dx = \left[\ln|x| + \frac{1}{x}\right]_1^2$

$= \left(\ln 2 + \frac{1}{2}\right) - (\ln 1 + 1)$

$= \ln 2 - \frac{1}{2} \approx 0.19$





14. By inspection, we see that the curves intersect at $x = 0$.

$A = \int_0^{\pi/2} (e^x - \cos x)\, dx = \left[e^x - \sin x\right]_0^{\pi/2}$

$= \left(e^{\pi/2} - \sin\frac{\pi}{2}\right) - (e^0 - \sin 0)$

$= [(e^{\pi/2} - 1) - (1 - 0)] = e^{\pi/2} - 2$





15. The curves intersect when $(x-2)^2 = x \;\Leftrightarrow\; x^2 - 4x + 4 = x \;\Leftrightarrow\; x^2 - 5x + 4 = 0 \;\Leftrightarrow\; (x-1)(x-4) = 0 \;\Leftrightarrow\; x = 1$ or 4.

$A = \int_1^4 [x - (x-2)^2]\, dx = \int_1^4 (-x^2 + 5x - 4)\, dx$

$= \left[-\frac{1}{3}x^3 + \frac{5}{2}x^2 - 4x\right]_1^4$

$= \left(-\frac{64}{3} + 40 - 16\right) - \left(-\frac{1}{3} + \frac{5}{2} - 4\right)$

$= \frac{9}{2}$





16. The curves intesect when $x^2 - 4x = 2x \;\Rightarrow\; x^2 - 6x = 0 \;\Rightarrow\; x(x-6) = 0 \;\Rightarrow\; x = 0$ or 6.

$A = \int_0^6 [2x - (x^2 - 4x)]\, dx$

$= \int_0^6 (6x - x^2)\, dx = \left[3x^2 - \frac{1}{3}x^3\right]_0^6$

$= \left[3(6)^2 - \frac{1}{3}(6)^3\right] - (0 - 0)$

$= 108 - 72 = 36$





17. The curves intersect when $1 - y^2 = y^2 - 1 \;\Leftrightarrow\; 2 = 2y^2 \;\Leftrightarrow\; y^2 = 1 \;\Leftrightarrow\; y = \pm 1$.

$A = \int_{-1}^{1} \left[(1 - y^2) - (y^2 - 1)\right] dy$

$= \int_{-1}^{1} 2(1 - y^2)\, dy = 2 \cdot 2 \int_0^1 (1 - y^2)\, dy$

$= 4\left[y - \frac{1}{3}y^3\right]_0^1 = 4\left(1 - \frac{1}{3}\right) = \frac{8}{3}$





© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

18. $4x + y^2 = 12$ and $x = y \quad \Rightarrow \quad 4x + x^2 = 12 \quad \Leftrightarrow \quad (x+6)(x-2) = 0 \quad \Leftrightarrow \quad x = -6$ or $x = 2$, so $y = -6$ or $y = 2$ and

$$\begin{aligned} A &= \int_{-6}^{2} \left[\left(-\tfrac{1}{4}y^2 + 3\right) - y \right] dy \\ &= \left[-\tfrac{1}{12}y^3 - \tfrac{1}{2}y^2 + 3y \right]_{-6}^{2} \\ &= \left(-\tfrac{2}{3} - 2 + 6\right) - (18 - 18 - 18) \\ &= 22 - \tfrac{2}{3} = \tfrac{64}{3} \end{aligned}$$





19. $12 - x^2 = x^2 - 6 \quad \Leftrightarrow \quad 2x^2 = 18 \quad \Leftrightarrow$
$x^2 = 9 \quad \Leftrightarrow \quad x = \pm 3$, so

$$\begin{aligned} A &= \int_{-3}^{3} \left[(12 - x^2) - (x^2 - 6) \right] dx \\ &= 2\int_{0}^{3} \left(18 - 2x^2\right) dx \qquad \text{[by symmetry]} \\ &= 2\left[18x - \tfrac{2}{3}x^3\right]_0^3 = 2\left[(54 - 18) - 0\right] \\ &= 2(36) = 72 \end{aligned}$$



20. $x^2 = 4x - x^2 \quad \Leftrightarrow \quad 2x^2 - 4x = 0 \quad \Leftrightarrow$
$2x(x-2) = 0 \quad \Leftrightarrow \quad x = 0$ or 2, so

$$\begin{aligned} A &= \int_{0}^{2} \left[(4x - x^2) - x^2 \right] dx = \int_{0}^{2} (4x - 2x^2)\, dx \\ &= \left[2x^2 - \tfrac{2}{3}x^3\right]_0^2 = 8 - \tfrac{16}{3} = \tfrac{8}{3} \end{aligned}$$



21. $2y^2 = 4 + y^2 \quad \Leftrightarrow \quad y^2 = 4 \quad \Leftrightarrow \quad y = \pm 2$, so

$$\begin{aligned} A &= \int_{-2}^{2} \left[(4 + y^2) - 2y^2 \right] dy \\ &= 2\int_{0}^{2} (4 - y^2)\, dy \qquad \text{[by symmetry]} \\ &= 2\left[4y - \tfrac{1}{3}y^3\right]_0^2 = 2\left(8 - \tfrac{8}{3}\right) = \tfrac{32}{3} \end{aligned}$$



22. The curves intersect when $\sqrt{x-1} = x - 1 \quad \Rightarrow$
$x - 1 = x^2 - 2x + 1 \quad \Leftrightarrow \quad 0 = x^2 - 3x + 2 \quad \Leftrightarrow$
$0 = (x-1)(x-2) \quad \Leftrightarrow \quad x = 1$ or 2.

$$\begin{aligned} A &= \int_{1}^{2} \left[\sqrt{x-1} - (x-1) \right] dx \\ &= \left[\tfrac{2}{3}(x-1)^{3/2} - \tfrac{1}{2}(x-1)^2 \right]_1^2 = \left(\tfrac{2}{3} - \tfrac{1}{2}\right) - (0 - 0) = \tfrac{1}{6} \end{aligned}$$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. $\sqrt[3]{2x} = \frac{1}{2}x \quad \Leftrightarrow \quad 2x = \left(\frac{1}{2}x\right)^3 = \frac{1}{8}x^3$

$\Leftrightarrow \quad 16x = x^3 \quad \Leftrightarrow \quad x^3 - 16x = 0$

$\Leftrightarrow \quad x(x^2 - 16) = 0 \quad \Leftrightarrow \quad x = -4,\ 0,\ \text{and } 4$



By symmetry,

$A = 2\int_0^4 \left(\sqrt[3]{2x} - \frac{1}{2}x\right) dx = 2\left[\frac{3}{8}(2x)^{4/3} - \frac{1}{4}x^2\right]_0^4$

$= 2[(6-4) - 0] = 4$

24. The curves intersect when $x^3 = x \quad \Leftrightarrow \quad x^3 - x = 0 \quad \Leftrightarrow$

$x(x^2-1) = 0 \quad \Leftrightarrow \quad x(x+1)(x-1) = 0 \quad \Leftrightarrow$

$x = 0$ or $x = \pm 1$.



$A = 2\int_0^1 (x - x^3)\, dx$ [by symmetry]

$= 2\left[\frac{1}{2}x^2 - \frac{1}{4}x^4\right]_0^1 = 2\left(\frac{1}{2} - \frac{1}{4}\right) = \frac{1}{2}$

25. $\sqrt{x} = \frac{1}{3}x \quad \Rightarrow \quad x = \frac{1}{9}x^2 \quad \Leftrightarrow \quad \frac{1}{9}x^2 - x = 0 \quad \Leftrightarrow$

$\frac{1}{9}x(x-9) = 0 \quad \Leftrightarrow \quad x = 0$ or $x = 9$



$A = \int_0^9 \left(\sqrt{x} - \frac{1}{3}x\right) dx + \int_9^{16} \left(\frac{1}{3}x - \sqrt{x}\right) dx$

$= \left[\frac{2}{3}x^{3/2} - \frac{1}{6}x^2\right]_0^9 + \left[\frac{1}{6}x^2 - \frac{2}{3}x^{3/2}\right]_9^{16}$

$= \left[\left(18 - \frac{27}{2}\right) - 0\right] + \left[\left(\frac{128}{3} - \frac{128}{3}\right) - \left(\frac{27}{2} - 18\right)\right] = 9$

26. $A = \int_0^{2\pi} [(2 - \cos x) - \cos x]\, dx$



$= \int_0^{2\pi} (2 - 2\cos x)\, dx$

$= \left[2x - 2\sin x\right]_0^{2\pi}$

$= (4\pi - 0) - 0 = 4\pi$

27. $\cos x = \sin 2x = 2\sin x \cos x \quad \Leftrightarrow \quad 2\sin x\cos x - \cos x = 0 \quad \Leftrightarrow \quad \cos x(2\sin x - 1) = 0 \quad \Leftrightarrow$

$\cos x = 0$ or $\sin x = \frac{1}{2} \quad \Leftrightarrow \quad x = \frac{\pi}{2}$ or $x = \frac{\pi}{6}$ on $\left[0, \frac{\pi}{2}\right]$



$A = \int_0^{\pi/6} (\cos x - \sin 2x)\, dx + \int_{\pi/6}^{\pi/2} (\sin 2x - \cos x)\, dx$

$= \left[\sin x + \frac{1}{2}\cos 2x\right]_0^{\pi/6} + \left[-\frac{1}{2}\cos 2x - \sin x\right]_{\pi/6}^{\pi/2}$

$= \left[\left(\sin\frac{\pi}{6} + \frac{1}{2}\cos\frac{\pi}{3}\right) - \left(\sin 0 + \frac{1}{2}\cos 0\right)\right]$

$\quad + \left[\left(-\frac{1}{2}\cos\pi - \sin\frac{\pi}{2}\right) - \left(-\frac{1}{2}\cos\frac{\pi}{3} - \sin\frac{\pi}{6}\right)\right]$

$= \left[\left(\frac{1}{2} + \frac{1}{4}\right) - \left(0 + \frac{1}{2}\right)\right] + \left[\left(\frac{1}{2} - 1\right) - \left(-\frac{1}{4} - \frac{1}{2}\right)\right] = \frac{1}{2}$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

28. The curves intersect when $\cos x = 1 - \cos x$ (on $[0, \pi]$) $\Leftrightarrow$ $2\cos x = 1$ $\Leftrightarrow$ $\cos x = \frac{1}{2}$ $\Leftrightarrow$ $x = \frac{\pi}{3}$.

$$\begin{aligned} A &= \int_0^{\pi/3} [\cos x - (1 - \cos x)]\,dx + \int_{\pi/3}^{\pi} [(1 - \cos x) - \cos x]\,dx \\ &= \int_0^{\pi/3} (2\cos x - 1)\,dx + \int_{\pi/3}^{\pi} (1 - 2\cos x)\,dx \\ &= \Big[2\sin x - x\Big]_0^{\pi/3} + \Big[x - 2\sin x\Big]_{\pi/3}^{\pi} \\ &= \left(\sqrt{3} - \frac{\pi}{3}\right) - 0 + (\pi - 0) - \left(\frac{\pi}{3} - \sqrt{3}\right) \\ &= 2\sqrt{3} + \frac{\pi}{3} \end{aligned}$$



29. The curves intersect when $8\cos x = \sec^2 x$ $\Rightarrow$ $8\cos^3 x = 1$ $\Rightarrow$ $\cos^3 x = \frac{1}{8}$ $\Rightarrow$ $\cos x = \frac{1}{2}$ $\Rightarrow$ $x = \frac{\pi}{3}$ for $0 < x < \frac{\pi}{2}$. By symmetry,

$$\begin{aligned} A &= 2\int_0^{\pi/3} (8\cos x - \sec^2 x)\,dx \\ &= 2\left[8\sin x - \tan x\right]_0^{\pi/3} \\ &= 2\left(8 \cdot \tfrac{\sqrt{3}}{2} - \sqrt{3}\right) = 2\left(3\sqrt{3}\right) \\ &= 6\sqrt{3} \end{aligned}$$





30. $x^4 - 3x^2 = x^2$ $\Leftrightarrow$ $x^4 - 4x^2 = 0$ $\Leftrightarrow$ $x^2(x^2 - 4) = 0$ $\Leftrightarrow$ $x = 0, x = \pm 2$

$$\begin{aligned} A &= 2\int_0^2 [x^2 - (x^4 - 3x^2)]\,dx = 2\int_0^2 (4x^2 - x^4)\,dx \\ &= 2\left[\frac{4}{3}x^3 - \frac{1}{5}x^5\right]_0^2 = 2\left[\left(\frac{32}{3} - \frac{32}{5}\right) - 0\right] \\ &= \frac{128}{15} \end{aligned}$$



31. By inspection, we see that the curves intersect at $x = \pm 1$ and that the area of the region enclosed by the curves is twice the area enclosed in the first quadrant.

$$\begin{aligned} A &= 2\int_0^1 [(2 - x) - x^4]\,dx = 2\left[2x - \tfrac{1}{2}x^2 - \tfrac{1}{5}x^5\right]_0^1 \\ &= 2\left[\left(2 - \tfrac{1}{2} - \tfrac{1}{5}\right) - 0\right] = 2\left(\tfrac{13}{10}\right) = \tfrac{13}{5} \end{aligned}$$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. $x^2 = \dfrac{32}{x^2+4} \quad \Leftrightarrow \quad x^2(x^2+4) = 32 \quad \Leftrightarrow$

$x^4 + 4x^2 - 32 = 0 \quad \Leftrightarrow \quad (x^2+8)(x^2-4) = 0 \quad \Leftrightarrow \quad x = \pm 2$



By symmetry,

$$A = 2\int_0^2 \left(\frac{32}{x^2+4} - x^2\right) dx$$

$$= 2\left[16\arctan\left(\frac{x}{2}\right) - \frac{1}{3}x^3\right]_0^2 = 2\left[\left(16\arctan 1 - \frac{8}{3}\right) - 0\right]$$

$$= 2\left(16\cdot\frac{\pi}{4} - \frac{8}{3}\right) = 8\pi - \frac{16}{3}$$

33. By inspection, we see that the curves intersect at $x = -1$, 0, and 1.



By symmetry,

$$A = 2\int_0^1 \left[\sin\left(\frac{\pi x}{2}\right) - x^3\right] dx = 2\left[-\frac{2}{\pi}\cos\left(\frac{\pi x}{2}\right) - \frac{1}{4}x^4\right]_0^1$$

$$= 2\left[\left(-\frac{2}{\pi}\cos\frac{\pi}{2} - \frac{1}{4}\right) - \left(-\frac{2}{\pi}\cos 0 - 0\right)\right]$$

$$= 2\left[\left(0 - \frac{1}{4}\right) - \left(-\frac{2}{\pi} - 0\right)\right] = 2\left(\frac{2}{\pi} - \frac{1}{4}\right) = \frac{4}{\pi} - \frac{1}{2}$$

34. $4 - 2\cosh x = \dfrac{1}{2}\sinh x \quad \Leftrightarrow \quad 4 - 2\left(\dfrac{e^x + e^{-x}}{2}\right) = \dfrac{1}{2}\left(\dfrac{e^x - e^{-x}}{2}\right) \quad \Leftrightarrow \quad 4 - e^x - e^{-x} = \dfrac{1}{4}e^x - \dfrac{1}{4}e^{-x} \quad \Leftrightarrow$

$4e^x(4 - e^x - e^{-x}) = 4e^x\left(\dfrac{1}{4}e^x - \dfrac{1}{4}e^{-x}\right) \quad \Leftrightarrow \quad 16e^x - 4e^{2x} - 4 = e^{2x} - 1 \quad \Leftrightarrow \quad 5e^{2x} - 16e^x + 3 = 0 \quad \Leftrightarrow$

$(5e^x - 1)(e^x - 3) = 0 \quad \Leftrightarrow \quad e^x = \frac{1}{5}$ or $e^x = 3 \quad \Leftrightarrow \quad x = \ln\frac{1}{5} = -\ln 5$ or $x = \ln 3$

$$A = \int_{-\ln 5}^{\ln 3} \left[(4 - 2\cosh x) - \frac{1}{2}\sinh x\right] dx = \left[4x - 2\sinh x - \frac{1}{2}\cosh x\right]_{-\ln 5}^{\ln 3}$$

$$= \left[4\ln 3 - 2\left(\frac{e^{\ln 3} - e^{-\ln 3}}{2}\right) - \frac{1}{2}\left(\frac{e^{\ln 3} + e^{-\ln 3}}{2}\right)\right] - \left[-4\ln 5 - 2\left(\frac{e^{-\ln 5} - e^{\ln 5}}{2}\right) - \frac{1}{2}\left(\frac{e^{-\ln 5} + e^{\ln 5}}{2}\right)\right]$$

$$= \left[4\ln 3 - \left(3 - \frac{1}{3}\right) - \frac{1}{4}\left(3 + \frac{1}{3}\right)\right] - \left[-4\ln 5 - \left(\frac{1}{5} - 5\right) - \frac{1}{4}\left(\frac{1}{5} + 5\right)\right]$$

$$= 4\ln 15 - 7$$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

35. $1/x = x \;\Leftrightarrow\; 1 = x^2 \;\Leftrightarrow\; x = \pm 1$ and $1/x = \frac{1}{4}x \;\Leftrightarrow$ $4 = x^2 \;\Leftrightarrow\; x = \pm 2$, so for $x > 0$,



$$\begin{aligned} A &= \int_0^1 \left(x - \frac{1}{4}x\right) dx + \int_1^2 \left(\frac{1}{x} - \frac{1}{4}x\right) dx \\ &= \int_0^1 \left(\frac{3}{4}x\right) dx + \int_1^2 \left(\frac{1}{x} - \frac{1}{4}x\right) dx \\ &= \left[\tfrac{3}{8}x^2\right]_0^1 + \left[\ln|x| - \tfrac{1}{8}x^2\right]_1^2 \\ &= \tfrac{3}{8} + \left(\ln 2 - \tfrac{1}{2}\right) - \left(0 - \tfrac{1}{8}\right) = \ln 2 \end{aligned}$$

36. $\frac{1}{4}x^2 = -x + 3 \;\Leftrightarrow\; x^2 + 4x - 12 = 0 \;\Leftrightarrow\; (x+6)(x-2) = 0 \;\Leftrightarrow\; x = -6$ or 2 and $2x^2 = -x + 3 \;\Leftrightarrow$ $2x^2 + x - 3 = 0 \;\Leftrightarrow\; (2x+3)(x-1) = 0 \;\Leftrightarrow\; x = -\frac{3}{2}$ or 1, so for $x \geq 0$,



$$\begin{aligned} A &= \int_0^1 \left(2x^2 - \tfrac{1}{4}x^2\right) dx + \int_1^2 \left[(-x+3) - \tfrac{1}{4}x^2\right] dx \\ &= \int_0^1 \tfrac{7}{4}x^2\, dx + \int_1^2 \left(-\tfrac{1}{4}x^2 - x + 3\right) dx \\ &= \left[\tfrac{7}{12}x^3\right]_0^1 + \left[-\tfrac{1}{12}x^3 - \tfrac{1}{2}x^2 + 3x\right]_1^2 \\ &= \tfrac{7}{12} + \left(-\tfrac{2}{3} - 2 + 6\right) - \left(-\tfrac{1}{12} - \tfrac{1}{2} + 3\right) = \tfrac{3}{2} \end{aligned}$$

37. (a) Total area $= 12 + 27 = 39$.

(b) $f(x) \leq g(x)$ for $0 \leq x \leq 2$ and $f(x) \geq g(x)$ for $2 \leq x \leq 5$, so

$$\begin{aligned} \textstyle\int_0^5 [f(x) - g(x)]\, dx &= \textstyle\int_0^2 [f(x) - g(x)]\, dx + \int_2^5 [f(x) - g(x)]\, dx \\ &= \textstyle -\int_0^2 [g(x) - f(x)]\, dx + \int_2^5 [f(x) - g(x)]\, dx \\ &= -(12) + 27 = 15 \end{aligned}$$

38. $\dfrac{x}{\sqrt{1+x^2}} = \dfrac{x}{\sqrt{9-x^2}} \;\Leftrightarrow\; x = 0$ or $\sqrt{1+x^2} = \sqrt{9-x^2} \;\Rightarrow$ $1 + x^2 = 9 - x^2 \;\Rightarrow\; 2x^2 = 8 \;\Rightarrow\; x^2 = 4 \;\Rightarrow\; x = 2$ $(x \geq 0)$.



$$\begin{aligned} A &= \int_0^2 \left(\frac{x}{\sqrt{1+x^2}} - \frac{x}{\sqrt{9-x^2}}\right) dx = \left[\sqrt{1+x^2} + \sqrt{9-x^2}\right]_0^2 \\ &= (\sqrt{5} + \sqrt{5}) - (1+3) = 2\sqrt{5} - 4 \end{aligned}$$

39. $\dfrac{x}{1+x^2} = \dfrac{x^2}{1+x^3} \;\Leftrightarrow\; x + x^4 = x^2 + x^4 \;\Leftrightarrow\; x = x^2 \;\Leftrightarrow$ $0 = x^2 - x \;\Leftrightarrow\; 0 = x(x-1) \;\Leftrightarrow\; x = 0$ or $x = 1$.



$$\begin{aligned} A &= \int_0^1 \left(\frac{x}{1+x^2} - \frac{x^2}{1+x^3}\right) dx = \left[\tfrac{1}{2}\ln(1+x^2) - \tfrac{1}{3}\ln(1+x^3)\right]_0^1 \\ &= \left(\tfrac{1}{2}\ln 2 - \tfrac{1}{3}\ln 2\right) - (0-0) = \tfrac{1}{6}\ln 2 \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. $\dfrac{\ln x}{x} = \dfrac{(\ln x)^2}{x} \quad \Leftrightarrow \quad \ln x = (\ln x)^2 \quad \Leftrightarrow \quad 0 = (\ln x)^2 - \ln x \quad \Leftrightarrow$
$0 = \ln x(\ln x - 1) \quad \Leftrightarrow \quad \ln x = 0 \text{ or } 1 \quad \Leftrightarrow \quad x = e^0 \text{ or } e^1 \ [1 \text{ or } e]$

$$A = \int_1^e \left[\frac{\ln x}{x} - \frac{(\ln x)^2}{x}\right] dx = \left[\tfrac{1}{2}(\ln x)^2 - \tfrac{1}{3}(\ln x)^3\right]_1^e$$
$$= \left(\tfrac{1}{2} - \tfrac{1}{3}\right) - (0 - 0) = \tfrac{1}{6}$$



41. An equation of the line through $(0, 0)$ and $(3, 1)$ is $y = \frac{1}{3}x$; through $(0, 0)$ and $(1, 2)$ is $y = 2x$; through $(3, 1)$ and $(1, 2)$ is $y = -\frac{1}{2}x + \frac{5}{2}$.

$$A = \int_0^1 \left(2x - \tfrac{1}{3}x\right) dx + \int_1^3 \left[\left(-\tfrac{1}{2}x + \tfrac{5}{2}\right) - \tfrac{1}{3}x\right] dx$$
$$= \int_0^1 \tfrac{5}{3}x\, dx + \int_1^3 \left(-\tfrac{5}{6}x + \tfrac{5}{2}\right) dx = \left[\tfrac{5}{6}x^2\right]_0^1 + \left[-\tfrac{5}{12}x^2 + \tfrac{5}{2}x\right]_1^3$$
$$= \tfrac{5}{6} + \left(-\tfrac{15}{4} + \tfrac{15}{2}\right) - \left(-\tfrac{5}{12} + \tfrac{5}{2}\right) = \tfrac{5}{2}$$



42. An equation of the line through $(2, 0)$ and $(0, 2)$ is $y = -x + 2$; through $(2, 0)$ and $(-1, 1)$ is $y = -\frac{1}{3}x + \frac{2}{3}$; through $(0, 2)$ and $(-1, 1)$ is $y = x + 2$.

$$A = \int_{-1}^0 \left[(x + 2) - \left(-\tfrac{1}{3}x + \tfrac{2}{3}\right)\right] dx + \int_0^2 \left[(-x + 2) - \left(-\tfrac{1}{3}x + \tfrac{2}{3}\right)\right] dx$$
$$= \int_{-1}^0 \left(\tfrac{4}{3}x + \tfrac{4}{3}\right) dx + \int_0^2 \left(-\tfrac{2}{3}x + \tfrac{4}{3}\right) dx$$
$$= \left[\tfrac{2}{3}x^2 + \tfrac{4}{3}x\right]_{-1}^0 + \left[-\tfrac{1}{3}x^2 + \tfrac{4}{3}x\right]_0^2$$
$$= 0 - \left(\tfrac{2}{3} - \tfrac{4}{3}\right) + \left(-\tfrac{4}{3} + \tfrac{8}{3}\right) - 0 = 2$$



43. The curves intersect when $\sin x = \cos 2x \quad (\text{on } [0, \pi/2]) \quad \Leftrightarrow \quad \sin x = 1 - 2\sin^2 x \quad \Leftrightarrow \quad 2\sin^2 x + \sin x - 1 = 0 \quad \Leftrightarrow$
$(2\sin x - 1)(\sin x + 1) = 0 \quad \Rightarrow \quad \sin x = \frac{1}{2} \quad \Rightarrow \quad x = \frac{\pi}{6}$.

$$A = \int_0^{\pi/2} |\sin x - \cos 2x|\, dx$$
$$= \int_0^{\pi/6} (\cos 2x - \sin x)\, dx + \int_{\pi/6}^{\pi/2} (\sin x - \cos 2x)\, dx$$
$$= \left[\tfrac{1}{2}\sin 2x + \cos x\right]_0^{\pi/6} + \left[-\cos x - \tfrac{1}{2}\sin 2x\right]_{\pi/6}^{\pi/2}$$
$$= \left(\tfrac{1}{4}\sqrt{3} + \tfrac{1}{2}\sqrt{3}\right) - (0 + 1) + (0 - 0) - \left(-\tfrac{1}{2}\sqrt{3} - \tfrac{1}{4}\sqrt{3}\right)$$
$$= \tfrac{3}{2}\sqrt{3} - 1$$



44. $$A = \int_{-1}^1 |3^x - 2^x|\, dx = \int_{-1}^0 (2^x - 3^x)\, dx + \int_0^1 (3^x - 2^x)\, dx$$
$$= \left[\frac{2^x}{\ln 2} - \frac{3^x}{\ln 3}\right]_{-1}^0 + \left[\frac{3^x}{\ln 3} - \frac{2^x}{\ln 2}\right]_0^1$$
$$= \left(\frac{1}{\ln 2} - \frac{1}{\ln 3}\right) - \left(\frac{1}{2\ln 2} - \frac{1}{3\ln 3}\right) + \left(\frac{3}{\ln 3} - \frac{2}{\ln 2}\right) - \left(\frac{1}{\ln 3} - \frac{1}{\ln 2}\right)$$
$$= \frac{2 - 1 - 4 + 2}{2\ln 2} + \frac{-3 + 1 + 9 - 3}{3\ln 3} = \frac{4}{3\ln 3} - \frac{1}{2\ln 2}$$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

45.



From the graph, we see that the curves intersect at $x = 0$ and $x = a \approx 0.896$, with $x\sin(x^2) > x^4$ on $(0, a)$. So the area A of the region bounded by the curves is

$$A = \int_0^a \left[x\sin(x^2) - x^4\right] dx = \left[-\tfrac{1}{2}\cos(x^2) - \tfrac{1}{5}x^5\right]_0^a$$
$$= -\tfrac{1}{2}\cos(a^2) - \tfrac{1}{5}a^5 + \tfrac{1}{2} \approx 0.037$$

46. From the graph, we see that the curves intersect (with $x \geq 0$) at $x = 0$ and $x = a$, where $a \approx 1.052$, with $x/(x^2+1)^2 > x^5 - x$ on $(0, a)$. The area A of the region bounded by the curves is

$$A = \int_0^a \left[\frac{x}{(x^2+1)^2} - (x^5 - x)\right] dx = \left[-\frac{1}{2}\cdot\frac{1}{x^2+1} - \frac{1}{6}x^6 + \frac{1}{2}x^2\right]_0^a$$
$$\approx 0.59$$



47.



From the graph, we see that the curves intersect at $x = a \approx -1.11$, $x = b \approx 1.25$, and $x = c \approx 2.86$, with $x^3 - 3x + 4 > 3x^2 - 2x$ on (a, b) and $3x^2 - 2x > x^3 - 3x + 4$ on (b, c). So the area of the region bounded by the curves is

$$A = \int_a^b \left[(x^3 - 3x + 4) - (3x^2 - 2x)\right] dx + \int_b^c \left[(3x^2 - 2x) - (x^3 - 3x + 4)\right] dx$$
$$= \int_a^b (x^3 - 3x^2 - x + 4)\, dx + \int_b^c (-x^3 + 3x^2 + x - 4)\, dx$$
$$= \left[\tfrac{1}{4}x^4 - x^3 - \tfrac{1}{2}x^2 + 4x\right]_a^b + \left[-\tfrac{1}{4}x^4 + x^3 + \tfrac{1}{2}x^2 - 4x\right]_b^c \approx 8.38$$

48.



From the graph, we see that the curves intersect at $x = a \approx 0.29$ and $x = b \approx 6.08$. $y = 2\sqrt{x}$ is the upper curve, so the area of the region bounded by the curves is

$$A \approx \int_a^b \left(2\sqrt{x} - 1.3^x\right) dx = \left[\tfrac{4}{3}x^{3/2} - \frac{1}{\ln 1.3}\,1.3^x\right]_a^b \approx 5.11$$

49.



Graph `Y1=2/(1+x^4)` and `Y2=x^2`. We see that `Y1` > `Y2` on $(-1, 1)$, so the area is given by $\int_{-1}^{1} \left(\frac{2}{1+x^4} - x^2\right) dx$. Evaluate the integral with a command such as `fnInt(Y1-Y2,x,-1,1)` to get 2.80123 to five decimal places.

Another method: Graph $f(x) =$ `Y1=2/(1+x^4)-x^2` and from the graph evaluate $\int f(x)\, dx$ from -1 to 1.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

50.



The curves intersect at $x = \pm 1$.

$$A = \int_{-1}^{1} (e^{1-x^2} - x^4)\, dx \approx 3.66016$$

51.



The curves intersect at $x = 0$ and $x = a \approx 0.749363$.

$$A = \int_{0}^{a} \left(\sqrt{x} - \tan^2 x\right)\, dx \approx 0.25142$$

52.



The curves intersect at $x = a \approx -1.911917$, $x = b \approx -1.223676$, and $x = c \approx 0.607946$.

$$A = \int_{a}^{b} \left[(x + 2\sin^4 x) - \cos x\right] dx + \int_{b}^{c} \left[\cos x - (x + 2\sin^4 x)\right] dx$$
$$\approx 1.70413$$

53. As the figure illustrates, the curves $y = x$ and $y = x^5 - 6x^3 + 4x$ enclose a four-part region symmetric about the origin (since $x^5 - 6x^3 + 4x$ and x are odd functions of x). The curves intersect at values of x where $x^5 - 6x^3 + 4x = x$; that is, where $x(x^4 - 6x^2 + 3) = 0$. That happens at $x = 0$ and where $x^2 = \dfrac{6 \pm \sqrt{36 - 12}}{2} = 3 \pm \sqrt{6}$; that is, at $x = -\sqrt{3 + \sqrt{6}}$, $-\sqrt{3 - \sqrt{6}}$, 0, $\sqrt{3 - \sqrt{6}}$, and $\sqrt{3 + \sqrt{6}}$. The exact area is



$$2\int_{0}^{\sqrt{3+\sqrt{6}}} \left|(x^5 - 6x^3 + 4x) - x\right| dx = 2\int_{0}^{\sqrt{3+\sqrt{6}}} \left|x^5 - 6x^3 + 3x\right| dx$$
$$= 2\int_{0}^{\sqrt{3-\sqrt{6}}} (x^5 - 6x^3 + 3x)\, dx + 2\int_{\sqrt{3-\sqrt{6}}}^{\sqrt{3+\sqrt{6}}} (-x^5 + 6x^3 - 3x)\, dx$$
$$\overset{\text{CAS}}{=} 12\sqrt{6} - 9$$

54. The inequality $x \geq 2y^2$ describes the region that lies on, or to the right of, the parabola $x = 2y^2$. The inequality $x \leq 1 - |y|$ describes the region that lies on, or to the left of, the curve $x = 1 - |y| = \begin{cases} 1 - y & \text{if } y \geq 0 \\ 1 + y & \text{if } y < 0 \end{cases}$. So the given region is the shaded region that lies between the curves.



[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

The graphs of $x = 1 - y$ and $x = 2y^2$ intersect when $1 - y = 2y^2 \quad \Leftrightarrow$

$2y^2 + y - 1 = 0 \quad \Leftrightarrow \quad (2y-1)(y+1) = 0 \quad \Rightarrow \quad y = \frac{1}{2}$ [for $y \geq 0$]. By symmetry,

$A = 2\int_0^{1/2} \left[(1-y) - 2y^2\right] dy = 2\left[-\frac{2}{3}y^3 - \frac{1}{2}y^2 + y\right]_0^{1/2} = 2\left[\left(-\frac{1}{12} - \frac{1}{8} + \frac{1}{2}\right) - 0\right] = 2\left(\frac{7}{24}\right) = \frac{7}{12}.$

55. 1 second $= \frac{1}{3600}$ hour, so 10 s $= \frac{1}{360}$ h. With the given data, we can take $n = 5$ to use the Midpoint Rule.

$\Delta t = \frac{1/360 - 0}{5} = \frac{1}{1800}$, so

$$\begin{aligned}
\text{distance}_{\text{Kelly}} - \text{distance}_{\text{Chris}} &= \int_0^{1/360} v_K\,dt - \int_0^{1/360} v_C\,dt = \int_0^{1/360} (v_K - v_C)\,dt \\
&\approx M_5 = \tfrac{1}{1800}\left[(v_K - v_C)(1) + (v_K - v_C)(3) + (v_K - v_C)(5)\right. \\
&\qquad \left. + (v_K - v_C)(7) + (v_K - v_C)(9)\right] \\
&= \tfrac{1}{1800}[(22-20) + (52-46) + (71-62) + (86-75) + (98-86)] \\
&= \tfrac{1}{1800}(2 + 6 + 9 + 11 + 12) = \tfrac{1}{1800}(40) = \tfrac{1}{45} \text{ mile, or } 117\tfrac{1}{3} \text{ feet}
\end{aligned}$$

56. If $x =$ distance from left end of pool and $w = w(x) =$ width at x, then the Midpoint Rule with $n = 4$ and $\Delta x = \dfrac{b-a}{n} = \dfrac{8 \cdot 2 - 0}{4} = 4$ gives Area $= \int_0^{16} w\,dx \approx 4(6.2 + 6.8 + 5.0 + 4.8) = 4(22.8) = 91.2 \text{ m}^2$.

57. Let $h(x)$ denote the height of the wing at x cm from the left end.

$$\begin{aligned}
A \approx M_5 &= \frac{200 - 0}{5}\left[h(20) + h(60) + h(100) + h(140) + h(180)\right] \\
&= 40(20.3 + 29.0 + 27.3 + 20.5 + 8.7) = 40(105.8) = 4232 \text{ cm}^2
\end{aligned}$$

58. For $0 \leq t \leq 10$, $b(t) > d(t)$, so the area between the curves is given by

$$\begin{aligned}
\int_0^{10} [b(t) - d(t)]\,dt &= \int_0^{10} (2200e^{0.024t} - 1460e^{0.018t})\,dt = \left[\frac{2200}{0.024}e^{0.024t} - \frac{1460}{0.018}e^{0.018t}\right]_0^{10} \\
&= \left(\frac{275{,}000}{3}e^{0.24} - \frac{730{,}000}{9}e^{0.18}\right) - \left(\frac{275{,}000}{3} - \frac{730{,}000}{9}\right) \approx 8868 \text{ people}
\end{aligned}$$

This area A represents the increase in population over a 10-year period.

59. (a) From Example 8(a), the infectiousness concentration is 1210 cells/mL. $g(t) = 1210 \quad \Leftrightarrow \quad 0.9f(t) = 1210 \quad \Leftrightarrow$ $0.9(-t)(t-21)(t+1) = 1210$. Using a calculator to solve the last equation for $t > 0$ gives us two solutions with the lesser being $t = t_3 \approx 11.26$ days, or the 12th day.

(b) From Example 8(b), the slope of the line through P_1 and P_2 is -23. From part (a), $P_3 = (t_3, 1210)$. An equation of the line through P_3 that is parallel to $\overline{P_1P_2}$ is $N - 1210 = -23(t - t_3)$, or $N = -23t + 23t_3 + 1210$. Using a calculator, we find that this line intersects g at $t = t_4 \approx 17.18$, or the 18th day. So in the patient with some immunity, the infection lasts about 2 days less than in the patient without immunity.

(c) The level of infectiousness for this patient is the area between the graph of g and the line in part (b). This area is

$$\begin{aligned}
\int_{t_3}^{t_4} \left[g(t) - (-23t + 23t_3 + 1210)\right] dt &\approx \int_{11.26}^{17.18} (-0.9t^3 + 18t^2 + 41.9t - 1468.94)\,dt \\
&= \left[-0.225t^4 + 6t^3 + 20.95t^2 - 1468.94t\right]_{11.26}^{17.18} \approx 706 \text{ (cells/mL)} \cdot \text{days}
\end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

60. From the figure, $g(t) > f(t)$ for $0 \le t \le 2$. The area between the curves is given by

$$\begin{aligned}\int_0^2 [g(t) - f(t)]\,dt &= \int_0^2 [(0.17t^2 - 0.5t + 1.1) - (0.73t^3 - 2t^2 + t + 0.6)]\,dt \\ &= \int_0^2 (-0.73t^3 + 2.17t^2 - 1.5t + 0.5)\,dt \\ &= \left[-\frac{0.73}{4}t^4 + \frac{2.17}{3}\,t^3 - 0.75t^2 + 0.5t\right]_0^2 \\ &= -2.92 + \frac{17.36}{3} - 3 + 1 - 0 = 0.8\overline{6} \approx 0.87\end{aligned}$$

Thus, about 0.87 more inches of rain fell at the second location than at the first during the first two hours of the storm.



61. We know that the area under curve A between $t = 0$ and $t = x$ is $\int_0^x v_A(t)\,dt = s_A(x)$, where $v_A(t)$ is the velocity of car A and s_A is its displacement. Similarly, the area under curve B between $t = 0$ and $t = x$ is $\int_0^x v_B(t)\,dt = s_B(x)$.

(a) After one minute, the area under curve A is greater than the area under curve B. So car A is ahead after one minute.

(b) The area of the shaded region has numerical value $s_A(1) - s_B(1)$, which is the distance by which A is ahead of B after 1 minute.

(c) After two minutes, car B is traveling faster than car A and has gained some ground, but the area under curve A from $t = 0$ to $t = 2$ is still greater than the corresponding area for curve B, so car A is still ahead.

(d) From the graph, it appears that the area between curves A and B for $0 \le t \le 1$ (when car A is going faster), which corresponds to the distance by which car A is ahead, seems to be about 3 squares. Therefore, the cars will be side by side at the time x where the area between the curves for $1 \le t \le x$ (when car B is going faster) is the same as the area for $0 \le t \le 1$. From the graph, it appears that this time is $x \approx 2.2$. So the cars are side by side when $t \approx 2.2$ minutes.

62. The area under $R'(x)$ from $x = 50$ to $x = 100$ represents the change in revenue, and the area under $C'(x)$ from $x = 50$ to $x = 100$ represents the change in cost. The shaded region represents the difference between these two values; that is, the increase in profit as the production level increases from 50 units to 100 units. We use the Midpoint Rule with $n = 5$ and $\Delta x = 10$:

$$\begin{aligned}M_5 &= \Delta x\{[R'(55) - C'(55)] + [R'(65) - C'(65)] + [R'(75) - C'(75)] + [R'(85) - C'(85)] + [R'(95) - C'(95)]\} \\ &\approx 10(2.40 - 0.85 + 2.20 - 0.90 + 2.00 - 1.00 + 1.80 - 1.10 + 1.70 - 1.20) \\ &= 10(5.05) = 50.5 \text{ thousand dollars}\end{aligned}$$

Using M_1 would give us $50(2 - 1) = 50$ thousand dollars.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

63.



To graph this function, we must first express it as a combination of explicit functions of y; namely, $y = \pm x\sqrt{x+3}$. We can see from the graph that the loop extends from $x = -3$ to $x = 0$, and that by symmetry, the area we seek is just twice the area under the top half of the curve on this interval, the equation of the top half being $y = -x\sqrt{x+3}$. So the area is $A = 2\int_{-3}^{0}\left(-x\sqrt{x+3}\right)dx$. We substitute $u = x+3$, so $du = dx$ and the limits change to 0 and 3, and we get

$$A = -2\int_0^3 [(u-3)\sqrt{u}\,]\,du = -2\int_0^3 (u^{3/2} - 3u^{1/2})\,du$$
$$= -2\left[\tfrac{2}{5}u^{5/2} - 2u^{3/2}\right]_0^3 = -2\left[\tfrac{2}{5}\left(3^2\sqrt{3}\right) - 2\left(3\sqrt{3}\right)\right] = \tfrac{24}{5}\sqrt{3}$$

64.



We start by finding the equation of the tangent line to $y = x^2$ at the point $(1, 1)$: $y' = 2x$, so the slope of the tangent is $2(1) = 2$, and its equation is $y - 1 = 2(x-1)$, or $y = 2x - 1$. We would need two integrals to integrate with respect to x, but only one to integrate with respect to y.

$$A = \int_0^1 \left[\tfrac{1}{2}(y+1) - \sqrt{y}\,\right]dy = \left[\tfrac{1}{4}y^2 + \tfrac{1}{2}y - \tfrac{2}{3}y^{3/2}\right]_0^1$$
$$= \tfrac{1}{4} + \tfrac{1}{2} - \tfrac{2}{3} = \tfrac{1}{12}$$

65.



By the symmetry of the problem, we consider only the first quadrant, where $y = x^2 \;\Rightarrow\; x = \sqrt{y}$. We are looking for a number b such that

$$\int_0^b \sqrt{y}\,dy = \int_b^4 \sqrt{y}\,dy \;\Rightarrow\; \tfrac{2}{3}\left[y^{3/2}\right]_0^b = \tfrac{2}{3}\left[y^{3/2}\right]_b^4 \;\Rightarrow$$
$$b^{3/2} = 4^{3/2} - b^{3/2} \;\Rightarrow\; 2b^{3/2} = 8 \;\Rightarrow\; b^{3/2} = 4 \;\Rightarrow\; b = 4^{2/3} \approx 2.52.$$

66. (a) We want to choose a so that

$$\int_1^a \frac{1}{x^2}\,dx = \int_a^4 \frac{1}{x^2}\,dx \;\Rightarrow\; \left[\frac{-1}{x}\right]_1^a = \left[\frac{-1}{x}\right]_a^4 \;\Rightarrow\; -\frac{1}{a} + 1 = -\frac{1}{4} + \frac{1}{a} \;\Rightarrow\; \frac{5}{4} = \frac{2}{a} \;\Rightarrow\; a = \frac{8}{5}.$$

(b) The area under the curve $y = 1/x^2$ from $x = 1$ to $x = 4$ is $\frac{3}{4}$ [take $a = 4$ in the first integral in part (a)]. Now the line $y = b$ must intersect the curve $x = 1/\sqrt{y}$ and not the line $x = 4$, since the area under the line $y = 1/4^2$ from $x = 1$ to $x = 4$ is only $\frac{3}{16}$, which is less than half of $\frac{3}{4}$. We want to choose b so that the upper area in the diagram is half of the total area under the curve $y = 1/x^2$ from $x = 1$ to $x = 4$. This implies that

$\int_b^1 \left(1/\sqrt{y} - 1\right)dy = \frac{1}{2}\cdot\frac{3}{4} \;\Rightarrow\; \left[2\sqrt{y} - y\right]_b^1 = \frac{3}{8} \;\Rightarrow\; 1 - 2\sqrt{b} + b = \frac{3}{8} \;\Rightarrow$
$b - 2\sqrt{b} + \frac{5}{8} = 0$. Letting $c = \sqrt{b}$, we get $c^2 - 2c + \frac{5}{8} = 0 \;\Rightarrow$
$8c^2 - 16c + 5 = 0$. Thus, $c = \frac{16 \pm \sqrt{256 - 160}}{16} = 1 \pm \frac{\sqrt{6}}{4}$. But $c = \sqrt{b} < 1 \;\Rightarrow$
$c = 1 - \frac{\sqrt{6}}{4} \;\Rightarrow\; b = c^2 = 1 + \frac{3}{8} - \frac{\sqrt{6}}{2} = \frac{1}{8}\left(11 - 4\sqrt{6}\right) \approx 0.1503.$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

67. We first assume that $c > 0$, since c can be replaced by $-c$ in both equations without changing the graphs, and if $c = 0$ the curves do not enclose a region. We see from the graph that the enclosed area A lies between $x = -c$ and $x = c$, and by symmetry, it is equal to four times the area in the first quadrant. The enclosed area is

$A = 4\int_0^c (c^2 - x^2)\,dx = 4\left[c^2x - \frac{1}{3}x^3\right]_0^c = 4\left(c^3 - \frac{1}{3}c^3\right) = 4\left(\frac{2}{3}c^3\right) = \frac{8}{3}c^3$

So $A = 576 \quad \Leftrightarrow \quad \frac{8}{3}c^3 = 576 \quad \Leftrightarrow \quad c^3 = 216 \quad \Leftrightarrow \quad c = \sqrt[3]{216} = 6.$

Note that $c = -6$ is another solution, since the graphs are the same.



68.



It appears from the diagram that the curves $y = \cos x$ and $y = \cos(x - c)$ intersect halfway between 0 and c, namely, when $x = c/2$. We can verify that this is indeed true by noting that $\cos(c/2 - c) = \cos(-c/2) = \cos(c/2)$. The point where $\cos(x - c)$ crosses the x-axis is $x = \frac{\pi}{2} + c$. So we require that $\int_0^{c/2} [\cos x - \cos(x - c)]\,dx = -\int_{\pi/2+c}^{\pi} \cos(x - c)\,dx$ [the negative sign on the RHS is needed since the second area is beneath the x-axis] $\Leftrightarrow$ $[\sin x - \sin(x - c)]_0^{c/2} = -[\sin(x - c)]_{\pi/2+c}^{\pi} \Rightarrow$ $[\sin(c/2) - \sin(-c/2)] - [-\sin(-c)] = -\sin(\pi - c) + \sin\left[\left(\frac{\pi}{2} + c\right) - c\right] \Leftrightarrow 2\sin(c/2) - \sin c = -\sin c + 1.$ [Here we have used the oddness of the sine function, and the fact that $\sin(\pi - c) = \sin c$]. So $2\sin(c/2) = 1 \Leftrightarrow$ $\sin(c/2) = \frac{1}{2} \quad \Leftrightarrow \quad c/2 = \frac{\pi}{6} \quad \Leftrightarrow \quad c = \frac{\pi}{3}$.

69. Let a and b be the x-coordinates of the points where the line intersects the curve. From the figure, $R_1 = R_2 \quad \Rightarrow$



$$\int_0^a \left[c - \left(8x - 27x^3\right)\right] dx = \int_a^b \left[\left(8x - 27x^3\right) - c\right] dx$$

$$\left[cx - 4x^2 + \tfrac{27}{4}x^4\right]_0^a = \left[4x^2 - \tfrac{27}{4}x^4 - cx\right]_a^b$$

$$ac - 4a^2 + \tfrac{27}{4}a^4 = \left(4b^2 - \tfrac{27}{4}b^4 - bc\right) - \left(4a^2 - \tfrac{27}{4}a^4 - ac\right)$$

$$0 = 4b^2 - \tfrac{27}{4}b^4 - bc = 4b^2 - \tfrac{27}{4}b^4 - b\left(8b - 27b^3\right)$$

$$= 4b^2 - \tfrac{27}{4}b^4 - 8b^2 + 27b^4 = \tfrac{81}{4}b^4 - 4b^2$$

$$= b^2\left(\tfrac{81}{4}b^2 - 4\right)$$

So for $b > 0$, $b^2 = \frac{16}{81} \quad \Rightarrow \quad b = \frac{4}{9}$. Thus, $c = 8b - 27b^3 = 8\left(\frac{4}{9}\right) - 27\left(\frac{64}{729}\right) = \frac{32}{9} - \frac{64}{27} = \frac{32}{27}$.

70. The curve and the line will determine a region when they intersect at two or more points. So we solve the equation $x/(x^2 + 1) = mx \quad \Rightarrow$

$x = x(mx^2 + m) \quad \Rightarrow \quad x(mx^2 + m) - x = 0 \quad \Rightarrow$

$x(mx^2 + m - 1) = 0 \quad \Rightarrow \quad x = 0$ or $mx^2 + m - 1 = 0 \quad \Rightarrow$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$x = 0$ or $x^2 = \dfrac{1-m}{m} \quad \Rightarrow \quad x = 0$ or $x = \pm\sqrt{\dfrac{1}{m} - 1}$. Note that if $m = 1$, this has only the solution $x = 0$, and no region is determined. But if $1/m - 1 > 0 \quad \Leftrightarrow \quad 1/m > 1 \quad \Leftrightarrow \quad 0 < m < 1$, then there are two solutions. [Another way of seeing this is to observe that the slope of the tangent to $y = x/(x^2+1)$ at the origin is $y'(0) = 1$ and therefore we must have $0 < m < 1$.] Note that we cannot just integrate between the positive and negative roots, since the curve and the line cross at the origin. Since mx and $x/(x^2+1)$ are both odd functions, the total area is twice the area between the curves on the interval $\left[0, \sqrt{1/m - 1}\right]$. So the total area enclosed is

$$2\int_0^{\sqrt{1/m-1}} \left[\frac{x}{x^2+1} - mx\right] dx = 2\left[\tfrac{1}{2}\ln(x^2+1) - \tfrac{1}{2}mx^2\right]_0^{\sqrt{1/m-1}} = [\ln(1/m - 1 + 1) - m(1/m - 1)] - (\ln 1 - 0)$$
$$= \ln(1/m) - 1 + m = m - \ln m - 1$$

APPLIED PROJECT The Gini Index

1. (a) $G = \dfrac{\text{area between } L \text{ and } y = x}{\text{area under } y = x} = \dfrac{\int_0^1 [x - L(x)]\,dx}{\frac{1}{2}} = 2\displaystyle\int_0^1 [x - L(x)]\,dx$

(b) For a perfectly egalitarian society, $L(x) = x$, so $G = 2\int_0^1 [x - x]\,dx = 0$. For a perfectly totalitarian society,

$$L(x) = \begin{cases} 1 & \text{if } x = 1 \\ 0 & \text{if } 0 \le x < 1 \end{cases} \quad \text{so } G = 2\int_0^1 (x - 0)\,dx = 2\left[\tfrac{1}{2}x^2\right]_0^1 = 2\left(\tfrac{1}{2}\right) = 1.$$

2. (a) The richest 20% of the population in 2016 received $1 - L(0.8) = 1 - 0.485 = 0.515$, or 51.5%, of the total US income.

(b) A quadratic model has the form $Q(x) = ax^2 + bx + c$. Rounding to six decimal places, we get $a = 1.341\,071$, $b = -0.411\,929$, and $c = 0.028\,571$. The quadratic model appears to be a reasonable fit, but note that $Q(0) \neq 0$ and Q is both decreasing and increasing.



(c) $G = 2\displaystyle\int_0^1 [x - Q(x)]\,dx \approx 0.4607$

3.

	$Q(x) = ax^2 + bx + c$			
Year	a	b	c	Gini
1980	1.149 554	−0.189 696	0.016 179	0.3910
1990	1.214 732	−0.265 589	0.020 393	0.4150
2000	1.280 804	−0.345 232	0.025 821	0.4397
2010	1.312 946	−0.378 518	0.026 679	0.4499

The Gini index has risen steadily from 1980 to 2016. The trend is toward a less egalitarian society.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. Using TI's `PwrReg` command and omitting the point $(0, 0)$ gives us $P(x) = 0.839\,312x^{2.103\,154}$ and a Gini index of $G = 2\int_0^1 [x - P(x)]\,dx \approx 0.4591$. Note that the power function is nearly quadratic.



6.2 Volumes

1. (a)



(b) A cross-section is a disk with radius $x^2 + 5$, so its area is

$$A(x) = \pi(x^2 + 5)^2 = \pi(x^4 + 10x^2 + 25).$$

$$V = \int_0^3 A(x)\,dx = \int_0^3 \pi(x^4 + 10x^2 + 25)\,dx$$

(c) $\displaystyle\int_0^3 \pi(x^4 + 10x^2 + 25)\,dx = \pi\left[\frac{1}{5}x^5 + \frac{10}{3}x^3 + 25x\right]_0^3 = \pi\left(\frac{243}{5} + 90 + 75\right) = \frac{1068}{5}\pi$

2. (a)



(b) A cross-section is a washer (annulus) with inner radius $\frac{1}{2}x$ and outer radius $\sqrt{x}$, so its area is

$$A(x) = \pi\left[\left(\sqrt{x}\,\right)^2 - \left(\tfrac{1}{2}x\right)^2\right] = \pi\left(x - \tfrac{1}{4}x^2\right).$$

$$V = \int_0^4 A(x)\,dx = \int_0^4 \pi\left(x - \tfrac{1}{4}x^2\right)dx$$

(c) $\displaystyle\int_0^4 \pi\left(x - \tfrac{1}{4}x^2\right)dx = \pi\left[\tfrac{1}{2}x^2 - \tfrac{1}{12}x^3\right]_0^4 = \pi\left(8 - \frac{16}{3}\right) = \frac{8}{3}\pi$

3. (a)



(b) $y = x^3 + 1 \;\Rightarrow\; y - 1 = x^3 \;\Rightarrow\; x = \sqrt[3]{y - 1}$. Therefore, a cross-section is a disk with radius $\sqrt[3]{y - 1}$, so its area is

$$A(y) = \pi\left(\sqrt[3]{y - 1}\,\right)^2 = \pi(y - 1)^{2/3}.$$

$$V = \int_1^9 A(y)\,dy = \int_1^9 \pi(y - 1)^{2/3}\,dy$$

(c) $\displaystyle\int_1^9 \pi(y - 1)^{2/3}\,dy = \pi\left[\frac{3}{5}(y - 1)^{5/3}\right]_1^9 = \frac{3}{5}\pi(32 - 0) = \frac{96}{5}\pi$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. (a)



(b) $y = x/2 \;\Rightarrow\; x = 2y$; $y = 2/x \;\Rightarrow\; x = 2/y$. A cross-section is a washer (annulus) with inner radius $2/y$ and outer radius $2y$, so its area is $A(y) = \pi\left[(2y)^2 - \left(\dfrac{2}{y}\right)^2\right] = \pi\left(4y^2 - \dfrac{4}{y^2}\right)$.

$$V = \int_1^2 A(y)\,dy = \int_1^2 \pi\left(4y^2 - \frac{4}{y^2}\right) dy$$

(c) $\displaystyle\int_1^2 \pi\left(4y^2 - \frac{4}{y^2}\right) dy = \pi\left[\frac{4}{3}y^3 + \frac{4}{y}\right]_1^2 = \pi\left[\left(\frac{32}{3} + 2\right) - \left(\frac{4}{3} + 4\right)\right] = \frac{22}{3}\pi$

5. $V = \displaystyle\int_1^3 \pi(\ln x)^2\,dx$



6. $V = \displaystyle\int_0^5 \pi\left(\sqrt{5-y}\right)^2 dy$



7. $8y = x^2 \;\Rightarrow\; x = \sqrt{8y}$ for $x \geq 0$; $y = \sqrt{x} \;\Rightarrow\; x = y^2$ for $y \geq 0$.

$y^2 = \sqrt{8y} \;\Rightarrow\; y^4 = 8y \;\Leftrightarrow\; y^4 - 8y = 0 \;\Leftrightarrow\; y(y^3 - 8) = 0 \;\Leftrightarrow$ $y = 0$ or $y = 2$.

$$V = \int_0^2 \pi\left[\left(\sqrt{8y}\right)^2 - (y^2)^2\right] dy$$



8. $(x-2)^2 = x + 10 \;\Rightarrow\; x^2 - 4x + 4 = x + 10 \;\Rightarrow$ $x^2 - 5x - 6 = 0 \;\Rightarrow\; (x+1)(x-6) = 0 \;\Rightarrow$ $x = -1$ or $x = 6$.

$$V = \int_{-1}^{6} \pi\{(x+10)^2 - [(x-2)^2]^2\}\,dx$$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

9. $V = \pi \int_0^{\pi} \left\{ [\sin x - (-2)]^2 - [0 - (-2)]^2 \right\} dx$

$= \pi \int_0^{\pi} \left[(\sin x + 2)^2 - 2^2 \right] dx$



10. $y = \sqrt{x} \quad \Rightarrow \quad x = y^2$ for $y \geq 0$.

$V = \int_0^2 \pi[(6 - y^2)^2 - (6 - 4)^2]\, dy$



11. A cross-section is a disk with radius $x + 1$, so its area is

$A(x) = \pi(x + 1)^2 = \pi(x^2 + 2x + 1)$.

$V = \int_0^2 A(x)\, dx = \int_0^2 \pi(x^2 + 2x + 1)\, dx$

$= \pi \left[\frac{1}{3}x^3 + x^2 + x \right]_0^2$

$= \pi\left(\frac{8}{3} + 4 + 2 \right) = \frac{26\pi}{3}$



12. A cross-section is a disk with radius $\dfrac{1}{x}$, so its area is $A(x) = \pi\left(\dfrac{1}{x}\right)^2 = \pi x^{-2}$.

$V = \int_1^4 A(x)\, dx = \int_1^4 \pi x^{-2}\, dx$

$= \pi \left[-x^{-1} \right]_1^4 = \pi\left(-\frac{1}{4} + 1 \right)$

$= \frac{3\pi}{4}$



13. A cross-section is a disk with radius $\sqrt{x - 1}$, so its area is $A(x) = \pi\left(\sqrt{x - 1}\right)^2 = \pi(x - 1)$.

$$V = \int_1^5 A(x)\, dx = \int_1^5 \pi(x - 1)\, dx = \pi \left[\tfrac{1}{2}x^2 - x \right]_1^5 = \pi\left[\left(\tfrac{25}{2} - 5 \right) - \left(\tfrac{1}{2} - 1 \right) \right] = 8\pi$$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. A cross-section is a disk with radius e^x, so its area is $A(x) = \pi(e^x)^2 = \pi e^{2x}$.

$$V = \int_{-1}^{1} A(x)\,dx = \int_{-1}^{1} \pi e^{2x}\,dx$$
$$= \pi\left[\tfrac{1}{2}e^{2x}\right]_{-1}^{1} = \tfrac{\pi}{2}(e^2 - e^{-2})$$





15. A cross-section is a disk with radius $2\sqrt{y}$, so its area is $A(y) = \pi\left(2\sqrt{y}\right)^2$.

$$V = \int_0^9 A(y)\,dy = \int_0^9 \pi\left(2\sqrt{y}\right)^2 dy = 4\pi\int_0^9 y\,dy$$
$$= 4\pi\left[\tfrac{1}{2}y^2\right]_0^9 = 2\pi(81) = 162\pi$$





16. A cross-section is a disk with radius $\frac{1}{2}y^2$, so its area is $A(y) = \pi\left(\frac{1}{2}y^2\right)^2 = \frac{1}{4}\pi y^4$.

$$V = \int_0^4 A(y)\,dy = \int_0^4 \pi\left(\tfrac{1}{4}y^4\right) dy$$
$$= \tfrac{\pi}{4}\left[\tfrac{1}{5}y^5\right]_0^4 = \tfrac{\pi}{20}(4^5) = \tfrac{256\pi}{5}$$





17. A cross-section is a washer with inner radius $\frac{1}{2}y$ and outer radius $\sqrt{y}$, so its area is

$$A(y) = \pi\left[\left(\sqrt{y}\right)^2 - \left(\tfrac{1}{2}y\right)^2\right] = \pi\left(y - \tfrac{1}{4}y^2\right).$$

$$V = \int_0^4 \pi\left(y - \tfrac{1}{4}y^2\right) dy = \pi\left[\tfrac{1}{2}y^2 - \tfrac{1}{12}y^3\right]_0^4$$
$$= \pi\left[\left(8 - \tfrac{16}{3}\right) - 0\right] = \tfrac{8}{3}\pi$$





18. A cross-section is a washer (annulus) with inner radius 2 and outer radius $6 - x^2$, so its area is $A(x) = \pi[(6 - x^2)^2 - 2^2] = \pi(x^4 - 12x^2 + 32)$.

$$V = \int_{-2}^{2} A(x)\,dx = 2\int_0^2 \pi(x^4 - 12x^2 + 32)\,dx$$
$$= 2\pi\left[\tfrac{1}{5}x^5 - 4x^3 + 32x\right]_0^2$$
$$= 2\pi\left(\tfrac{32}{5} - 32 + 64\right) = 2\pi\left(\tfrac{192}{5}\right) = \tfrac{384\pi}{5}$$





© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

19. A cross-section is a washer with inner radius x^3 and outer radius $\sqrt{x}$, so its area is

$$A(x) = \pi\left[\left(\sqrt{x}\right)^2 - (x^3)^2\right] = \pi(x - x^6).$$

$$V = \int_0^1 \pi(x - x^6)\,dx = \pi\left[\tfrac{1}{2}x^2 - \tfrac{1}{7}x^7\right]_0^1$$
$$= \pi\left[\left(\tfrac{1}{2} - \tfrac{1}{7}\right) - 0\right] = \tfrac{5}{14}\pi$$





20. A cross-section is a washer with inner radius y^4 and outer radius $2 - y^2$, so its area is

$$A(y) = \pi(2-y^2)^2 - \pi(y^4)^2 = \pi(4 - 4y^2 + y^4 - y^8).$$

$$V = \int_{-1}^{1} A(y)\,dy = 2\int_0^1 \pi(4 - 4y^2 + y^4 - y^8)\,dy$$
$$= 2\pi\left[4y - \tfrac{4}{3}y^3 + \tfrac{1}{5}y^5 - \tfrac{1}{9}y^8\right]_0^1$$
$$= 2\pi\left(4 - \tfrac{4}{3} + \tfrac{1}{5} - \tfrac{1}{9}\right) = 2\pi\left(\tfrac{124}{45}\right) = \tfrac{248\pi}{45}$$





21. A cross-section is a washer with inner radius $1 - \sqrt{x}$ and outer radius $1 - x^2$, so its area is

$$A(x) = \pi\left[(1 - x^2)^2 - (1 - \sqrt{x}\,)^2\right]$$
$$= \pi\left[(1 - 2x^2 + x^4) - (1 - 2\sqrt{x} + x)\right]$$
$$= \pi\left(x^4 - 2x^2 + 2\sqrt{x} - x\right).$$

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi(x^4 - 2x^2 + 2x^{1/2} - x)\,dx$$
$$= \pi\left[\tfrac{1}{5}x^5 - \tfrac{2}{3}x^3 + \tfrac{4}{3}x^{3/2} - \tfrac{1}{2}x^2\right]_0^1$$
$$= \pi\left(\tfrac{1}{5} - \tfrac{2}{3} + \tfrac{4}{3} - \tfrac{1}{2}\right) = \tfrac{11}{30}\pi$$





22. A cross-section is a washer with inner radius $1 - (-3) = 4$ and outer radius $x^3 - (-3) = x^3 + 3$, so its area is

$$A(x) = \pi(x^3 + 3)^2 - \pi(4)^2 = \pi(x^6 + 6x^3 - 7).$$

$$V = \int_1^2 A(x)\,dx = \int_1^2 \pi(x^6 + 6x^3 - 7)\,dx$$
$$= \pi\left[\tfrac{1}{7}x^7 + \tfrac{3}{2}x^4 - 7x\right]_1^2$$
$$= \pi\left[\left(\tfrac{128}{7} + 24 - 14\right) - \left(\tfrac{1}{7} + \tfrac{3}{2} - 7\right)\right] = \tfrac{471\pi}{14}$$





© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

23. A cross-section is a washer with inner radius $(1+\sec x)-1=\sec x$ and outer radius $3-1=2$, so its area is

$$A(x)=\pi\left[2^2-(\sec x)^2\right]=\pi(4-\sec^2 x).$$

$$V=\int_{-\pi/3}^{\pi/3}A(x)\,dx=\int_{-\pi/3}^{\pi/3}\pi(4-\sec^2 x)\,dx$$

$$=2\pi\int_0^{\pi/3}(4-\sec^2 x)\,dx \qquad \text{[by symmetry]}$$

$$=2\pi\Big[4x-\tan x\Big]_0^{\pi/3}=2\pi\left[\left(\tfrac{4\pi}{3}-\sqrt{3}\right)-0\right]$$

$$=2\pi\left(\tfrac{4\pi}{3}-\sqrt{3}\right)$$





24. A cross-section is a washer with inner radius $\sin x-(-1)$ and outer radius $\cos x-(-1)$, so its area is

$$A(x)=\pi\left[(\cos x+1)^2-(\sin x+1)^2\right]$$

$$=\pi(\cos^2 x+2\cos x-\sin^2 x-2\sin x)$$

$$=\pi(\cos 2x+2\cos x-2\sin x).$$

$$V=\int_0^{\pi/4}A(x)\,dx=\int_0^{\pi/4}\pi(\cos 2x+2\cos x-2\sin x)\,dx$$

$$=\pi\left[\tfrac{1}{2}\sin 2x+2\sin x+2\cos x\right]_0^{\pi/4}$$

$$=\pi\left[\left(\tfrac{1}{2}+\sqrt{2}+\sqrt{2}\right)-(0+0+2)\right]=\left(2\sqrt{2}-\tfrac{3}{2}\right)\pi$$





25. A cross-section is a washer with inner radius $2-1$ and outer radius $2-\sqrt[3]{y}$, so its area is

$$A(y)=\pi\left[(2-\sqrt[3]{y})^2-(2-1)^2\right]=\pi\left[4-4\sqrt[3]{y}+\sqrt[3]{y^2}-1\right].$$

$$V=\int_0^1 A(y)\,dy=\int_0^1\pi(3-4y^{1/3}+y^{2/3})\,dy=\pi\left[3y-3y^{4/3}+\tfrac{3}{5}y^{2/3}\right]_0^1=\pi\left(3-3+\tfrac{3}{5}\right)=\tfrac{3}{5}\pi.$$





26. For $0\le y<\frac{1}{2}$, a cross-section is a washer with inner radius $1-(-1)$ and outer radius $2-(-1)$, so its area is $A(y)=\pi(3^2-2^2)=5\pi$. For $\frac{1}{2}\le y\le 1$, a cross-section is a washer with inner radius $1-(-1)$ and outer radius $1/y-(-1)$, so its area is $A(y)=\pi\left[(1/y+1)^2-(2)^2\right]=\pi(1/y^2+2/y+1-4)$.





[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$V = \int_0^{1/2} 5\pi\,dy + \int_{1/2}^{1} \pi\left(\frac{1}{y^2} + \frac{2}{y} - 3\right) dy = 5\pi\left[y\right]_0^{1/2} + \pi\left[-\frac{1}{y} + 2\ln y - 3y\right]_{1/2}^{1}$$
$$= 5\pi\left(\tfrac{1}{2} - 0\right) + \pi\left[(-1 + 0 - 3) - \left(-2 + 2\ln\tfrac{1}{2} - \tfrac{3}{2}\right)\right] = \tfrac{5}{2}\pi + \pi\left(-\tfrac{1}{2} + 2\ln 2\right)$$
$$= (2 + 2\ln 2)\pi = 2\pi(1 + \ln 2)$$

27. From the symmetry of the curves, we see they intersect at $x = \frac{1}{2}$ and so $y^2 = \frac{1}{2} \quad \Leftrightarrow \quad y = \pm\sqrt{\frac{1}{2}}$. A cross-section is a washer with inner radius $3 - (1 - y^2)$ and outer radius $3 - y^2$, so its area is

$$A(y) = \pi\left[(3 - y^2)^2 - (2 + y^2)^2\right]$$
$$= \pi\left[(9 - 6y^2 + y^4) - (4 + 4y^2 + y^4)\right]$$
$$= \pi(5 - 10y^2).$$

$$V = \int_{-\sqrt{1/2}}^{\sqrt{1/2}} A(y)\,dy$$
$$= 2\int_0^{\sqrt{1/2}} 5\pi(1 - 2y^2)\,dy \qquad \text{[by symmetry]}$$
$$= 10\pi\left[y - \tfrac{2}{3}y^3\right]_0^{\sqrt{2}/2} = 10\pi\left(\tfrac{\sqrt{2}}{2} - \tfrac{\sqrt{2}}{6}\right)$$
$$= 10\pi\left(\tfrac{\sqrt{2}}{3}\right) = \tfrac{10}{3}\sqrt{2}\,\pi$$



28. For $0 \le y < 2$, a cross-section is an annulus with inner radius $2 - 1$ and outer radius $4 - 1$, the area of which is $A_1(y) = \pi(4 - 1)^2 - \pi(2 - 1)^2$. For $2 \le y \le 4$, a cross-section is an annulus with inner radius $y - 1$ and outer radius $4 - 1$, the area of which is $A_2(y) = \pi(4 - 1)^2 - \pi(y - 1)^2$.

$$V = \int_0^4 A(y)\,dy = \pi\int_0^2 \left[(4 - 1)^2 - (2 - 1)^2\right] dy + \pi\int_2^4 \left[(4 - 1)^2 - (y - 1)^2\right] dy$$
$$= \pi\left[\,8y\,\right]_0^2 + \pi\int_2^4 (8 + 2y - y^2)\,dy$$
$$= 16\pi + \pi\left[8y + y^2 - \tfrac{1}{3}y^3\right]_2^4$$
$$= 16\pi + \pi\left[\left(32 + 16 - \tfrac{64}{3}\right) - \left(16 + 4 - \tfrac{8}{3}\right)\right]$$
$$= \tfrac{76}{3}\pi$$



29. $\mathcal{R}_1$ about OA (the line $y = 0$):

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi(x)^2\,dx = \pi\left[\tfrac{1}{3}x^3\right]_0^1 = \tfrac{1}{3}\pi$$

30. $\mathcal{R}_1$ about OC (the line $x = 0$):

$$V = \int_0^1 A(y)\,dy = \int_0^1 \pi(1^2 - y^2)\,dy = \pi\left[y - \tfrac{1}{3}y^3\right]_0^1 = \pi\left(1 - \tfrac{1}{3}\right) = \tfrac{2}{3}\pi$$

31. $\mathcal{R}_1$ about AB (the line $x = 1$):

$$V = \int_0^1 A(y)\,dy = \int_0^1 \pi(1 - y)^2\,dy = \pi\int_0^1 (1 - 2y + y^2)\,dy = \pi\left[y - y^2 + \tfrac{1}{3}y^3\right]_0^1 = \tfrac{1}{3}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. $\mathcal{R}_1$ about BC (the line $y = 1$):

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi[(1-0)^2 - (1-x)^2]\,dx = \pi\int_0^1 [1-(1-2x+x^2)]\,dx$$
$$= \pi\int_0^1 (-x^2+2x)\,dx = \pi\left[-\tfrac{1}{3}x^3 + x^2\right]_0^1 = \pi\left(-\tfrac{1}{3}+1\right) = \tfrac{2}{3}\pi$$

33. $\mathcal{R}_2$ about OA (the line $y = 0$):

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi\left[1^2 - \left(\sqrt[4]{x}\right)^2\right]dx = \pi\int_0^1 (1-x^{1/2})\,dx = \pi\left[x - \tfrac{2}{3}x^{3/2}\right]_0^1 = \pi\left(1-\tfrac{2}{3}\right) = \tfrac{1}{3}\pi$$

34. $\mathcal{R}_2$ about OC (the line $x = 0$):

$$V = \int_0^1 A(y)\,dy = \int_0^1 \pi[(y^4)^2]\,dy = \pi\int_0^1 y^8\,dy = \pi\left[\tfrac{1}{9}y^9\right]_0^1 = \tfrac{1}{9}\pi$$

35. $\mathcal{R}_2$ about AB (the line $x = 1$):

$$V = \int_0^1 A(y)\,dy = \int_0^1 \pi[1^2 - (1-y^4)^2]\,dy = \pi\int_0^1 [1-(1-2y^4+y^8)]\,dy$$
$$= \pi\int_0^1 (2y^4 - y^8)\,dy = \pi\left[\tfrac{2}{5}y^5 - \tfrac{1}{9}y^9\right]_0^1 = \pi\left(\tfrac{2}{5}-\tfrac{1}{9}\right) = \tfrac{13}{45}\pi$$

36. $\mathcal{R}_2$ about BC (the line $y = 1$):

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi(1-\sqrt[4]{x})^2\,dx = \pi\int_0^1 (1-2x^{1/4}+x^{1/2})\,dx$$
$$= \pi\left[x - \tfrac{8}{5}x^{5/4} + \tfrac{2}{3}x^{3/2}\right]_0^1 = \pi\left(1-\tfrac{8}{5}+\tfrac{2}{3}\right) = \tfrac{1}{15}\pi$$

37. $\mathcal{R}_3$ about OA (the line $y = 0$):

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi\left[\left(\sqrt[4]{x}\right)^2 - x^2\right]dx = \pi\int_0^1 (x^{1/2}-x^2)\,dx = \pi\left[\tfrac{2}{3}x^{3/2} - \tfrac{1}{3}x^3\right]_0^1 = \pi\left(\tfrac{2}{3}-\tfrac{1}{3}\right) = \tfrac{1}{3}\pi$$

Note: Let $\mathcal{R} = \mathcal{R}_1 \cup \mathcal{R}_2 \cup \mathcal{R}_3$. If we rotate $\mathcal{R}$ about any of the segments OA, OC, AB, or BC, we obtain a right circular cylinder of height 1 and radius 1. Its volume is $\pi r^2 h = \pi(1)^2 \cdot 1 = \pi$. As a check for Exercises 29, 33, and 37, we can add the answers, and that sum must equal π. Thus, $\frac{1}{3}\pi + \frac{1}{3}\pi + \frac{1}{3}\pi = \pi$.

38. $\mathcal{R}_3$ about OC (the line $x = 0$):

$$V = \int_0^1 A(y)\,dy = \int_0^1 \pi[y^2 - (y^4)^2]\,dy = \pi\int_0^1 (y^2-y^8)\,dy = \pi\left[\tfrac{1}{3}y^3 - \tfrac{1}{9}y^9\right]_0^1 = \pi\left(\tfrac{1}{3}-\tfrac{1}{9}\right) = \tfrac{2}{9}\pi$$

Note: See the note in the solution to Exercise 37. For Exercises 30, 34, and 38, we have $\frac{2}{3}\pi + \frac{1}{9}\pi + \frac{2}{9}\pi = \pi$.

39. $\mathcal{R}_3$ about AB (the line $x = 1$):

$$V = \int_0^1 A(y)\,dy = \int_0^1 \pi[(1-y^4)^2 - (1-y)^2]\,dy = \pi\int_0^1 [(1-2y^4+y^8) - (1-2y+y^2)]\,dy$$
$$= \pi\int_0^1 (y^8 - 2y^4 - y^2 + 2y)\,dy = \pi\left[\tfrac{1}{9}y^9 - \tfrac{2}{5}y^5 - \tfrac{1}{3}y^3 + y^2\right]_0^1 = \pi\left(\tfrac{1}{9}-\tfrac{2}{5}-\tfrac{1}{3}+1\right) = \tfrac{17}{45}\pi$$

Note: See the note in the solution to Exercise 37. For Exercises 31, 35, and 39, we have $\frac{1}{3}\pi + \frac{13}{45}\pi + \frac{17}{45}\pi = \pi$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

40. $\mathcal{R}_3$ about BC (the line $y = 1$):

$$V = \int_0^1 A(x)\,dx = \int_0^1 \pi[(1-x)^2 - \left(1 - \sqrt[4]{x}\right)^2]\,dx = \pi \int_0^1 [(1 - 2x + x^2) - (1 - 2x^{1/4} + x^{1/2})]\,dx$$

$$= \pi \int_0^1 (x^2 - 2x - x^{1/2} + 2x^{1/4})\,dx = \pi\left[\tfrac{1}{3}x^3 - x^2 - \tfrac{2}{3}x^{3/2} + \tfrac{8}{5}x^{5/4}\right]_0^1 = \pi\left(\tfrac{1}{3} - 1 - \tfrac{2}{3} + \tfrac{8}{5}\right) = \tfrac{4}{15}\pi$$

Note: See the note in the solution to Exercise 37. For Exercises 32, 36, and 40, we have $\frac{2}{3}\pi + \frac{1}{15}\pi + \frac{4}{15}\pi = \pi$.

41. (a) About the x-axis:



$$V = \int_{-1}^1 \pi(e^{-x^2})^2\,dx = 2\pi \int_0^1 e^{-2x^2}\,dx \quad \text{[by symmetry]}$$

$$\approx 3.75825$$

(b) About $y = -1$:



$$V = \int_{-1}^1 \pi\left\{[e^{-x^2} - (-1)]^2 - [0 - (-1)]^2\right\}\,dx$$

$$= 2\pi \int_0^1 [(e^{-x^2} + 1)^2 - 1]\,dx = 2\pi \int_0^1 (e^{-2x^2} + 2e^{-x^2})\,dx$$

$$\approx 13.14312$$

42. (a) About the x-axis:



$$V = \int_{-\pi/2}^{\pi/2} \pi(\cos^2 x)^2\,dx = 2\pi \int_0^{\pi/2} \cos^4 x\,dx \quad \text{[by symmetry]}$$

$$\approx 3.70110$$

(b) About $y = 1$:



$$V = \int_{-\pi/2}^{\pi/2} \pi[(1-0)^2 - (1 - \cos^2 x)^2]\,dx$$

$$= 2\pi \int_0^{\pi/2} [1 - (1 - 2\cos^2 x + \cos^4 x)]\,dx$$

$$= 2\pi \int_0^{\pi/2} (2\cos^2 x - \cos^4 x)\,dx \approx 6.16850$$

43. (a) About $y = 2$:



$$x^2 + 4y^2 = 4 \quad \Rightarrow \quad 4y^2 = 4 - x^2 \quad \Rightarrow \quad y^2 = 1 - x^2/4 \quad \Rightarrow$$

$$y = \pm\sqrt{1 - x^2/4}$$

$$V = \int_{-2}^2 \pi\left\{\left[2 - \left(-\sqrt{1 - x^2/4}\right)\right]^2 - \left(2 - \sqrt{1 - x^2/4}\right)^2\right\}\,dx$$

$$= 2\pi \int_0^2 8\sqrt{1 - x^2/4}\,dx \approx 78.95684$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) About $x = 2$:

$$x^2 + 4y^2 = 4 \quad \Rightarrow \quad x^2 = 4 - 4y^2 \quad \Rightarrow \quad x = \pm\sqrt{4 - 4y^2}$$

$$V = \int_{-1}^{1} \pi \left\{ \left[2 - \left(-\sqrt{4 - 4y^2} \right) \right]^2 - \left(2 - \sqrt{4 - 4y^2} \right)^2 \right\} dy$$

$$= 2\pi \int_0^1 8\sqrt{4 - 4y^2}\, dy \approx 78.95684$$



[Notice that this is the same approximation as in part (a). This can be explained by Pappus's Theorem in Section 8.3.]

44. (a) About the x-axis:

$$y = x^2 \text{ and } x^2 + y^2 = 1 \quad \Rightarrow \quad x^2 + x^4 = 1 \quad \Rightarrow \quad x^4 + x^2 - 1 = 0 \quad \Rightarrow$$

$$x^2 = \frac{-1 + \sqrt{5}}{2} \approx 0.618 \quad \Rightarrow \quad x = \pm a = \pm\sqrt{\frac{-1 + \sqrt{5}}{2}} \approx \pm 0.786.$$

$$V = \int_{-a}^{a} \pi \left[\left(\sqrt{1 - x^2} \right)^2 - (x^2)^2 \right] dx = 2\pi \int_0^a (1 - x^2 - x^4)\, dx$$

$$\approx 3.54459$$



(b) About the y-axis:

$$V = \int_0^{a^2} \pi \left(\sqrt{y}\right)^2 dy + \int_{a^2}^1 \pi \left(\sqrt{1 - y^2} \right)^2 dy$$

$$= \pi \int_0^{a^2} y\, dy + \pi \int_{a^2}^1 (1 - y^2)\, dy \approx 0.99998$$



45. $y = \ln(x^6 + 2)$ and $y = \sqrt{3 - x^3}$ intersect at $x = a \approx -4.091$, $x = b \approx -1.467$, and $x = c \approx 1.091$.



$$V = \pi \int_a^b \left\{ \left[\ln(x^6 + 2) \right]^2 - \left(\sqrt{3 - x^3} \right)^2 \right\} dx + \pi \int_b^c \left\{ \left(\sqrt{3 - x^3} \right)^2 - \left[\ln(x^6 + 2) \right]^2 \right\} dx \approx 89.023$$

46. $y = 1 + xe^{-x^3}$ and $y = \arctan x^2$ intersect at $x = a \approx -0.570$ and $x = b \approx 1.391$.

$$V = \pi \int_a^b \left[\left(1 + xe^{-x^3} \right)^2 - (\arctan x^2)^2 \right] dx \approx 6.923$$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. $V = \pi \int_0^{\pi} \left\{ \left[\sin^2 x - (-1)\right]^2 - \left[0 - (-1)\right]^2 \right\} dx$

$\overset{\text{CAS}}{=} \frac{11}{8}\pi^2$



48. $V = \pi \int_0^2 \left[(3-x)^2 - (3 - xe^{1-x/2})^2\right] dx$

$\overset{\text{CAS}}{=} \pi\left(-2e^2 + 24e - \frac{142}{3}\right)$



49. $\pi \int_0^{\pi/2} \sin^2 x\, dx = \pi \int_0^{\pi/2} (\sin x)^2\, dx$ describes the volume of the solid obtained by rotating the region $\mathcal{R} = \{(x, y) \mid 0 \le x \le \pi/2, 0 \le y \le \sin x\}$ of the xy-plane about the x-axis.

50. $\pi \int_0^{\ln 2} e^{2x}\, dx = \pi \int_0^{\ln 2} (e^x)^2\, dx$ describes the volume of the solid obtained by rotating the region $\mathcal{R} = \{(x, y) \mid 0 \le x \le \ln 2, 0 \le y \le e^x\}$ of the xy-plane about the x-axis.

51. $\pi \int_0^1 (x^4 - x^6)\, dx = \pi \int_0^1 [(x^2)^2 - (x^3)^2]\, dx$ describes the volume of the solid obtained by rotating the region $\mathcal{R} = \{(x, y) \mid 0 \le x \le 1, x^3 \le y \le x^2\}$ of the xy-plane about the x-axis.

52. $\pi \int_{-1}^1 (1 - y^2)^2\, dy$ describes the volume of the solid obtained by rotating the region $\mathcal{R} = \{(x, y) \mid -1 \le y \le 1, 0 \le x \le 1 - y^2\}$ of the xy-plane about the y-axis.

53. $\pi \int_0^4 y\, dy = \pi \int_0^4 \left(\sqrt{y}\right)^2 dy$ describes the volume of the solid obtained by rotating the region $\mathcal{R} = \left\{(x, y) \mid 0 \le y \le 4, 0 \le x \le \sqrt{y}\right\}$ of the xy-plane about the y-axis.

54. $\pi \int_1^4 \left[3^2 - \left(3 - \sqrt{x}\right)^2\right] dx$ describes the volume of the solid obtained by rotating the region $\mathcal{R} = \left\{(x, y) \mid 1 \le x \le 4, 3 - \sqrt{x} \le y \le 3\right\}$ of the xy-plane about the x-axis.

55. There are 10 subintervals over the 15-cm length, so we'll use $n = 10/2 = 5$ for the Midpoint Rule.

$$V = \int_0^{15} A(x)\, dx \approx M_5 = \tfrac{15-0}{5}[A(1.5) + A(4.5) + A(7.5) + A(10.5) + A(13.5)]$$
$$= 3(18 + 79 + 106 + 128 + 39) = 3 \cdot 370 = 1110 \text{ cm}^3$$

56. $V = \int_0^{10} A(x)\, dx \approx M_5 = \tfrac{10-0}{5}[A(1) + A(3) + A(5) + A(7) + A(9)]$

$$= 2(0.65 + 0.61 + 0.59 + 0.55 + 0.50) = 2(2.90) = 5.80 \text{ m}^3$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

57. (a) $V = \int_2^{10} \pi\,[f(x)]^2\,dx \approx \pi\frac{10-2}{4}\left\{[f(3)]^2 + [f(5)]^2 + [f(7)]^2 + [f(9)]^2\right\}$

$\approx 2\pi\left[(1.5)^2 + (2.2)^2 + (3.8)^2 + (3.1)^2\right] \approx 196 \text{ units}^3$

(b) $V = \int_0^4 \pi\left[(\text{outer radius})^2 - (\text{inner radius})^2\right] dy$

$\approx \pi\frac{4-0}{4}\left\{\left[(9.9)^2 - (2.2)^2\right] + \left[(9.7)^2 - (3.0)^2\right] + \left[(9.3)^2 - (5.6)^2\right] + \left[(8.7)^2 - (6.5)^2\right]\right\}$

$\approx 838 \text{ units}^3$

58. (a) $V = \displaystyle\int_{-1}^{1} \pi\left[\left(ax^3 + bx^2 + cx + d\right)\sqrt{1-x^2}\right]^2 dx \overset{\text{CAS}}{=} \frac{4\left\{5a^2 + 18ac + 3\left[3b^2 + 14bd + 7(c^2 + 5d^2)\right]\right\}\pi}{315}$

(b) $y = (-0.06x^3 + 0.04x^2 + 0.1x + 0.54)\sqrt{1-x^2}$ is graphed in the figure. Substitute $a = -0.06$, $b = 0.04$, $c = 0.1$, and $d = 0.54$ in the answer for part (a) to get $V \overset{\text{CAS}}{=} \dfrac{3769\pi}{9375} \approx 1.263$.



59. We'll form a right circular cone with height h and base radius r by revolving the line $y = \frac{r}{h}x$ about the x-axis.



$$V = \pi\int_0^h \left(\frac{r}{h}\,x\right)^2 dx = \pi\int_0^h \frac{r^2}{h^2}\,x^2 dx = \pi\,\frac{r^2}{h^2}\left[\frac{1}{3}x^3\right]_0^h$$

$$= \pi\,\frac{r^2}{h^2}\left(\frac{1}{3}h^3\right) = \frac{1}{3}\pi r^2 h$$

Another solution: Revolve $x = -\dfrac{r}{h}\,y + r$ about the y-axis.



$$V = \pi\int_0^h \left(-\frac{r}{h}y + r\right)^2 dy \overset{*}{=} \pi\int_0^h \left[\frac{r^2}{h^2}y^2 - \frac{2r^2}{h}y + r^2\right] dy$$

$$= \pi\left[\frac{r^2}{3h^2}y^3 - \frac{r^2}{h}y^2 + r^2 y\right]_0^h = \pi\left(\tfrac{1}{3}r^2h - r^2h + r^2h\right) = \tfrac{1}{3}\pi r^2 h$$

* Or use substitution with $u = r - \dfrac{r}{h}\,y$ and $du = -\dfrac{r}{h}\,dy$ to get

$$\pi\int_r^0 u^2\left(-\frac{h}{r}\,du\right) = -\pi\,\frac{h}{r}\left[\frac{1}{3}u^3\right]_r^0 = -\pi\,\frac{h}{r}\left(-\frac{1}{3}r^3\right) = \frac{1}{3}\pi r^2 h.$$

60. $V = \displaystyle\pi\int_0^h \left(R - \frac{R-r}{h}\,y\right)^2 dy$

$$= \pi\int_0^h \left[R^2 - \frac{2R(R-r)}{h}\,y + \left(\frac{R-r}{h}\right)^2 y^2\right] dy$$

$$= \pi\left[R^2 y - \frac{R(R-r)}{h}\,y^2 + \frac{1}{3}\left(\frac{R-r}{h}\right)^2 y^3\right]_0^h$$

$$= \pi\left[R^2h - R(R-r)h + \tfrac{1}{3}(R-r)^2 h\right]$$

$$= \tfrac{1}{3}\pi h\left[3Rr + (R^2 - 2Rr + r^2)\right] = \tfrac{1}{3}\pi h(R^2 + Rr + r^2)$$



[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Another solution: $\dfrac{H}{R}=\dfrac{H-h}{r}$ by similar triangles. Therefore, $Hr=HR-hR \;\Rightarrow\; hR=H(R-r) \;\Rightarrow$ $H=\dfrac{hR}{R-r}$. Now

$$\begin{aligned}V&=\tfrac{1}{3}\pi R^2H-\tfrac{1}{3}\pi r^2(H-h) \qquad \text{[by Exercise 59]}\\ &=\tfrac{1}{3}\pi R^2\,\frac{hR}{R-r}-\tfrac{1}{3}\pi r^2\frac{rh}{R-r} \qquad \left[H-h=\frac{rH}{R}=\frac{rhR}{R(R-r)}\right]\\ &=\frac{1}{3}\pi h\,\frac{R^3-r^3}{R-r}=\tfrac{1}{3}\pi h\big(R^2+Rr+r^2\big)\\ &=\tfrac{1}{3}\Big[\pi R^2+\pi r^2+\sqrt{(\pi R^2)\,(\pi r^2)}\,\Big]h=\tfrac{1}{3}\big(A_1+A_2+\sqrt{A_1A_2}\,\big)h\end{aligned}$$



where A_1 and A_2 are the areas of the bases of the frustum. (See Exercise 62 for a related result.)

61. $x^2+y^2=r^2 \;\Leftrightarrow\; x^2=r^2-y^2$

$$\begin{aligned}V&=\pi\int_{r-h}^{r}\big(r^2-y^2\big)\,dy=\pi\left[r^2y-\frac{y^3}{3}\right]_{r-h}^{r}=\pi\left\{\left[r^3-\frac{r^3}{3}\right]-\left[r^2(r-h)-\frac{(r-h)^3}{3}\right]\right\}\\ &=\pi\big\{\tfrac{2}{3}r^3-\tfrac{1}{3}(r-h)\big[3r^2-(r-h)^2\big]\big\}\\ &=\tfrac{1}{3}\pi\left\{2r^3-(r-h)\big[3r^2-\big(r^2-2rh+h^2\big)\big]\right\}\\ &=\tfrac{1}{3}\pi\left\{2r^3-(r-h)\big[2r^2+2rh-h^2\big]\right\}\\ &=\tfrac{1}{3}\pi\big(2r^3-2r^3-2r^2h+rh^2+2r^2h+2rh^2-h^3\big)\\ &=\tfrac{1}{3}\pi\big(3rh^2-h^3\big)=\tfrac{1}{3}\pi h^2(3r-h), \text{ or, equivalently, } \pi h^2\left(r-\frac{h}{3}\right)\end{aligned}$$



62. An equation of the line is $x=\dfrac{\Delta x}{\Delta y}\,y+(x\text{-intercept})=\dfrac{a/2-b/2}{h-0}\,y+\dfrac{b}{2}=\dfrac{a-b}{2h}\,y+\dfrac{b}{2}$.

$$\begin{aligned}V&=\int_0^h A(y)\,dy=\int_0^h(2x)^2\,dy\\ &=\int_0^h\left[2\left(\frac{a-b}{2h}y+\frac{b}{2}\right)\right]^2dy=\int_0^h\left[\frac{a-b}{h}y+b\right]^2dy\\ &=\int_0^h\left[\frac{(a-b)^2}{h^2}y^2+\frac{2b(a-b)}{h}y+b^2\right]dy\\ &=\left[\frac{(a-b)^2}{3h^2}y^3+\frac{b(a-b)}{h}y^2+b^2y\right]_0^h\\ &=\tfrac{1}{3}(a-b)^2h+b(a-b)h+b^2h=\tfrac{1}{3}\big(a^2-2ab+b^2+3ab\big)h\\ &=\tfrac{1}{3}\big(a^2+ab+b^2\big)h\end{aligned}$$



[Note that this can be written as $\tfrac{1}{3}\big(A_1+A_2+\sqrt{A_1A_2}\,\big)h$, as in Exercise 60.]

If $a=b$, we get a rectangular solid with volume b^2h. If $a=0$, we get a square pyramid with volume $\tfrac{1}{3}b^2h$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

63. For a cross-section at height y, we see from similar triangles that $\dfrac{\alpha/2}{b/2} = \dfrac{h-y}{h}$, so $\alpha = b\left(1-\dfrac{y}{h}\right)$.

Similarly, for cross-sections having $2b$ as their base and β replacing α, $\beta = 2b\left(1-\dfrac{y}{h}\right)$. So



$$V = \int_0^h A(y)\,dy = \int_0^h \left[b\left(1-\frac{y}{h}\right)\right]\left[2b\left(1-\frac{y}{h}\right)\right]dy$$
$$= \int_0^h 2b^2\left(1-\frac{y}{h}\right)^2 dy = 2b^2\int_0^h \left(1-\frac{2y}{h}+\frac{y^2}{h^2}\right)dy$$
$$= 2b^2\left[y-\frac{y^2}{h}+\frac{y^3}{3h^2}\right]_0^h = 2b^2\left[h-h+\tfrac{1}{3}h\right]$$
$$= \tfrac{2}{3}b^2h \quad [= \tfrac{1}{3}Bh \text{ where } B \text{ is the area of the base, as with any pyramid.}]$$

64. Consider the triangle consisting of two vertices of the base and the center of the base. This triangle is similar to the corresponding triangle at a height y, so $a/b = \alpha/\beta \;\Rightarrow\; \alpha = a\beta/b$. Also by similar triangles, $b/h = \beta/(h-y) \Rightarrow \beta = b(h-y)/h$. These two equations imply that $\alpha = a(1-y/h)$, and since the cross-section is an equilateral triangle, it has area



$$A(y) = \frac{1}{2}\cdot\alpha\cdot\frac{\sqrt{3}}{2}\alpha = \frac{a^2(1-y/h)^2}{4}\sqrt{3}, \text{ so}$$
$$V = \int_0^h A(y)\,dy = \frac{a^2\sqrt{3}}{4}\int_0^h \left(1-\frac{y}{h}\right)^2 dy$$
$$= \frac{a^2\sqrt{3}}{4}\left[-\frac{h}{3}\left(1-\frac{y}{h}\right)^3\right]_0^h = -\frac{\sqrt{3}}{12}a^2h(-1) = \frac{\sqrt{3}}{12}a^2h$$

65. A cross-section at height z is a triangle similar to the base, so we'll multiply the legs of the base triangle, 3 and 4, by a proportionality factor of $(5-z)/5$. Thus, the triangle at height z has area



$$A(z) = \frac{1}{2}\cdot 3\left(\frac{5-z}{5}\right)\cdot 4\left(\frac{5-z}{5}\right) = 6\left(1-\frac{z}{5}\right)^2, \text{ so}$$
$$V = \int_0^5 A(z)\,dz = 6\int_0^5 \left(1-\frac{z}{5}\right)^2 dz = 6\int_1^0 u^2(-5\,du) \qquad \begin{bmatrix} u = 1-z/5, \\ du = -\frac{1}{5}\,dz \end{bmatrix}$$
$$= -30\left[\tfrac{1}{3}u^3\right]_1^0 = -30\left(-\tfrac{1}{3}\right) = 10 \text{ cm}^3$$

66. A cross-section is shaded in the diagram.



$$A(x) = (2y)^2 = \left(2\sqrt{r^2-x^2}\right)^2, \text{ so}$$
$$V = \textstyle\int_{-r}^{r} A(x)\,dx = 2\int_0^r 4(r^2-x^2)\,dx$$
$$= 8\left[r^2x-\tfrac{1}{3}x^3\right]_0^r = 8\left(\tfrac{2}{3}r^3\right) = \tfrac{16}{3}r^3$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

67. If l is a leg of the isosceles right triangle and $2y$ is the hypotenuse,



then $l^2 + l^2 = (2y)^2 \quad \Rightarrow \quad 2l^2 = 4y^2 \quad \Rightarrow \quad l^2 = 2y^2$.

$$V = \int_{-2}^{2} A(x)\,dx = 2\int_0^2 A(x)\,dx = 2\int_0^2 \tfrac{1}{2}(l)(l)\,dx = 2\int_0^2 y^2\,dx$$
$$= 2\int_0^2 \tfrac{1}{4}(36 - 9x^2)\,dx = \tfrac{9}{2}\int_0^2 (4 - x^2)\,dx$$
$$= \tfrac{9}{2}\left[4x - \tfrac{1}{3}x^3\right]_0^2 = \tfrac{9}{2}\left(8 - \tfrac{8}{3}\right) = 24$$

68. The cross-section of the base corresponding to the coordinate y has length $x = 1 - y$. The corresponding equilateral triangle with side s has area $A(y) = s^2\left(\frac{\sqrt{3}}{4}\right) = (1-y)^2\left(\frac{\sqrt{3}}{4}\right)$. Therefore,



$$V = \int_0^1 A(y)\,dy = \int_0^1 (1-y)^2\left(\frac{\sqrt{3}}{4}\right)dy$$
$$= \frac{\sqrt{3}}{4}\int_0^1 (1 - 2y + y^2)\,dy = \frac{\sqrt{3}}{4}\left[y - y^2 + \tfrac{1}{3}y^3\right]_0^1$$
$$= \frac{\sqrt{3}}{4}\left(\frac{1}{3}\right) = \frac{\sqrt{3}}{12}$$

Or: $\displaystyle\int_0^1 (1-y)^2\left(\frac{\sqrt{3}}{4}\right)dy = \frac{\sqrt{3}}{4}\int_1^0 u^2(-du) \quad [u = 1-y] = \frac{\sqrt{3}}{4}\left[\frac{1}{3}u^3\right]_0^1 = \frac{\sqrt{3}}{12}$

69. The cross-section of the base corresponding to the coordinate x has length $y = 1 - x$. The corresponding square with side s has area $A(x) = s^2 = (1-x)^2 = 1 - 2x + x^2$. Therefore,



$$V = \int_0^1 A(x)\,dx = \int_0^1 (1 - 2x + x^2)\,dx$$
$$= \left[x - x^2 + \tfrac{1}{3}x^3\right]_0^1 = \left(1 - 1 + \tfrac{1}{3}\right) - 0 = \tfrac{1}{3}$$

Or: $\displaystyle\int_0^1 (1-x)^2\,dx = \int_1^0 u^2(-du) \quad [u = 1-x] = \left[\tfrac{1}{3}u^3\right]_0^1 = \tfrac{1}{3}$

70. The cross-section of the base corresponding to the coordinate y has length $2x = 2\sqrt{1-y}$. $\left[y = 1 - x^2 \quad \Leftrightarrow \quad x = \pm\sqrt{1-y}\right]$ The corresponding square with side s has area $A(x) = s^2 = \left(2\sqrt{1-y}\right)^2 = 4(1-y)$. Therefore,



$V = \int_0^1 A(y)\,dy = \int_0^1 4(1-y)\,dy = 4\left[y - \frac{1}{2}y^2\right]_0^1 = 4\left[\left(1 - \frac{1}{2}\right) - 0\right] = 2.$

71. The cross-section of the base b corresponding to the coordinate x has length $1 - x^2$. The height h also has length $1 - x^2$, so the corresponding isosceles triangle has area $A(x) = \frac{1}{2}bh = \frac{1}{2}(1-x^2)^2$. Therefore,



$$V = \int_{-1}^{1} A(x)\,dx = \int_{-1}^{1} \tfrac{1}{2}(1-x^2)^2\,dx$$
$$= 2\cdot\tfrac{1}{2}\int_0^1 (1 - 2x^2 + x^4)\,dx \qquad \text{[by symmetry]}$$
$$= \left[x - \tfrac{2}{3}x^3 + \tfrac{1}{5}x^5\right]_0^1 = \left(1 - \tfrac{2}{3} + \tfrac{1}{5}\right) - 0 = \tfrac{8}{15}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

72. The cross-section of the base corresponding to the coordinate y has length $2x = 2\sqrt{2-y}$. $[y = 2 - x^2 \quad \Leftrightarrow$ $x = \pm\sqrt{2-y}\,]$ The corresponding cross-section of the solid S is a quarter-circle with radius $2\sqrt{2-y}$ and area $A(y) = \frac{1}{4}\pi(2\sqrt{2-y})^2 = \pi(2-y)$. Therefore,



$$V = \int_0^2 A(y)\,dy = \int_0^2 \pi(2-y)\,dy$$
$$= \pi\left[2y - \tfrac{1}{2}y^2\right]_0^2 = \pi(4-2) = 2\pi$$

73. The cross-section of S at coordinate x, $-1 \le x \le 1$, is a circle centered at the point $\left(x, \frac{1}{2}(1-x^2)\right)$ with radius $\frac{1}{2}(1-x^2)$. The area of the cross-section is



$$A(x) = \pi\left[\tfrac{1}{2}(1-x^2)\right]^2 = \tfrac{\pi}{4}(1 - 2x^2 + x^4)$$

The volume of S is

$$V = \int_{-1}^{1} A(x)\,dx = 2\int_0^1 \tfrac{\pi}{4}(1 - 2x^2 + x^4)\,dx = \tfrac{\pi}{2}\left[x - \tfrac{2}{3}x^3 + \tfrac{1}{5}x^5\right]_0^1 = \tfrac{\pi}{2}\left(1 - \tfrac{2}{3} + \tfrac{1}{5}\right) = \tfrac{\pi}{2}\left(\tfrac{8}{15}\right) = \tfrac{4\pi}{15}$$

74. The cross-section of S at coordinate x, $0 \le x \le 4$, is a circle centered at the point $\left(x, \frac{1}{2}\left(\frac{1}{2}\sqrt{x} + \sqrt{x}\right)\right)$ with radius $\frac{1}{2}\left(\sqrt{x} - \frac{1}{2}\sqrt{x}\right)$. The area of the cross-section is $A(x) = \pi\left[\frac{1}{2}\left(\sqrt{x} - \frac{1}{2}\sqrt{x}\right)\right]^2 = \pi \cdot \frac{1}{4} \cdot \left(\frac{1}{2}\sqrt{x}\right)^2 = \dfrac{\pi x}{16}$. The volume of S is $V = \displaystyle\int_0^4 A(x)\,dx = \int_0^4 \frac{\pi x}{16}\,dx = \frac{\pi}{32}\left[x^2\right]_0^4 = \frac{\pi}{32}(16-0) = \frac{\pi}{2}$.

75. (a) The torus is obtained by rotating the circle $(x-R)^2 + y^2 = r^2$ about the y-axis. Solving for x, we see that the right half of the circle is given by $x = R + \sqrt{r^2 - y^2} = f(y)$ and the left half by $x = R - \sqrt{r^2 - y^2} = g(y)$. So



$$V = \pi\textstyle\int_{-r}^{r}\left\{[f(y)]^2 - [g(y)]^2\right\}dy$$
$$= 2\pi\textstyle\int_0^r\left[\left(R^2 + 2R\sqrt{r^2-y^2} + r^2 - y^2\right) - \left(R^2 - 2R\sqrt{r^2-y^2} + r^2 - y^2\right)\right]dy$$
$$= 2\pi\textstyle\int_0^r 4R\sqrt{r^2-y^2}\,dy = 8\pi R\int_0^r \sqrt{r^2-y^2}\,dy$$

(b) Observe that the integral represents a quarter of the area of a circle with radius r, so $8\pi R\int_0^r \sqrt{r^2-y^2}\,dy = 8\pi R \cdot \frac{1}{4}\pi r^2 = 2\pi^2 r^2 R$.

76. (a) $V = \int_{-r}^{r} A(x)\,dx = 2\int_0^r A(x)\,dx = 2\int_0^r \frac{1}{2}h\left(2\sqrt{r^2-x^2}\right)dx = 2h\int_0^r \sqrt{r^2-x^2}\,dx$

(b) Observe that the integral represents one quarter of the area of a circle of radius r, so $V = 2h \cdot \frac{1}{4}\pi r^2 = \frac{1}{2}\pi h r^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

77. The cross-sections perpendicular to the y-axis in Figure 17 are rectangles. The rectangle corresponding to the coordinate y has a base of length $2\sqrt{16-y^2}$ in the xy-plane and a height of $\frac{1}{\sqrt{3}}\,y$, since $\angle BAC = 30^\circ$ and $|BC| = \frac{1}{\sqrt{3}}\,|AB|$. Thus, $A(y) = \frac{2}{\sqrt{3}}\,y\sqrt{16-y^2}$ and

$$V = \int_0^4 A(y)\,dy = \tfrac{2}{\sqrt{3}}\int_0^4 \sqrt{16-y^2}\,y\,dy = \tfrac{2}{\sqrt{3}}\int_{16}^0 u^{1/2}\left(-\tfrac{1}{2}\,du\right) \qquad [\text{Put } u = 16-y^2, \text{ so } du = -2y\,dy]$$

$$= \tfrac{1}{\sqrt{3}}\int_0^{16} u^{1/2}\,du = \tfrac{1}{\sqrt{3}}\,\tfrac{2}{3}\left[u^{3/2}\right]_0^{16} = \tfrac{2}{3\sqrt{3}}\,(64) = \tfrac{128}{3\sqrt{3}}$$

78. (a) $\text{Volume}(S_1) = \int_0^h A(z)\,dz = \text{Volume}(S_2)$ since the cross-sectional area $A(z)$ at height z is the same for both solids.

(b) By Cavalieri's Principle, the volume of the cylinder in the figure is the same as that of a right circular cylinder with radius r and height h, that is, $\pi r^2 h$.

79.



By similar triangles, the radius x_1 of the cross-section at height y of the cone removed from the cylinder satisfies $\dfrac{x_1}{y} = \dfrac{r}{r} \quad \Rightarrow \quad x_1 = y$. Thus, the area of the annular cross-section at height y remaining once the cone is removed from the cylinder is $\pi(r^2 - y^2)$.

The radius x_2 of the cross-section at height y of the hemisphere satisfies $x_2^2 + y^2 = r^2 \quad \Rightarrow \quad x_2 = \sqrt{r^2-y^2}$.

The area of the circular cross-section at height y is then $\pi\left(\sqrt{r^2-y^2}\right)^2 = \pi(r^2-y^2)$.

Each cross-section at height y of the cylinder with cone removed has area equal to that of the corresponding cross-section at height y of the hemisphere. By Cavalieri's Principle, the volumes of the solids are then equal.

80. Each cross-section of the solid S in a plane perpendicular to the x-axis is a square (since the edges of the cut lie on the cylinders, which are perpendicular). One-quarter of this square and one-eighth of S are shown. The area of this quarter-square is $|PQ|^2 = r^2 - x^2$. Therefore, $A(x) = 4(r^2-x^2)$ and the volume of S is



$$V = \int_{-r}^{r} A(x)\,dx = 4\int_{-r}^{r}(r^2-x^2)\,dx$$

$$= 8(r^2-x^2)\,dx = 8\left[r^2x - \tfrac{1}{3}x^3\right]_0^r = \tfrac{16}{3}r^3$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

81. The volume is obtained by rotating the area common to two circles of radius r, as shown. The volume of the right half is



$$V_{\text{right}} = \pi \int_0^{r/2} y^2\,dx = \pi \int_0^{r/2} \left[r^2 - \left(\tfrac{1}{2}r + x\right)^2\right] dx$$

$$= \pi\left[r^2x - \tfrac{1}{3}\left(\tfrac{1}{2}r + x\right)^3\right]_0^{r/2} = \pi\left[\left(\tfrac{1}{2}r^3 - \tfrac{1}{3}r^3\right) - \left(0 - \tfrac{1}{24}r^3\right)\right] = \tfrac{5}{24}\pi r^3$$

So by symmetry, the total volume is twice this, or $\frac{5}{12}\pi r^3$.

Another solution: We observe that the volume is the twice the volume of a cap of a sphere, so we can use the formula from Exercise 61 with $h = \frac{1}{2}r$: $V = 2 \cdot \frac{1}{3}\pi h^2(3r - h) = \frac{2}{3}\pi\left(\frac{1}{2}r\right)^2\left(3r - \frac{1}{2}r\right) = \frac{5}{12}\pi r^3$.

82. We consider two cases: one in which the ball is not completely submerged and the other in which it is.

Case 1: $0 \le h \le 10$ The ball will not be completely submerged, and so a cross-section of the water parallel to the surface will be the shaded area shown in the first diagram. We can find the area of the cross-section at height x above the bottom of the bowl by using the Pythagorean Theorem: $R^2 = 15^2 - (15 - x)^2$ and $r^2 = 5^2 - (x - 5)^2$, so $A(x) = \pi\left(R^2 - r^2\right) = 20\pi x$. The volume of water when it has depth h is then $V(h) = \int_0^h A(x)\,dx = \int_0^h 20\pi x\,dx = \left[10\pi x^2\right]_0^h = 10\pi h^2$ cm^3, $0 \le h \le 10$.

Case 2: $10 < h \le 15$ In this case we can find the volume by simply subtracting the volume displaced by the ball from the total volume inside the bowl underneath the surface of the water. The total volume underneath the surface is just the volume of a cap of the bowl, so we use the formula from Exercise 61: $V_{\text{cap}}(h) = \frac{1}{3}\pi h^2(45 - h)$. The volume of the small sphere is $V_{\text{ball}} = \frac{4}{3}\pi(5)^3 = \frac{500}{3}\pi$, so the total volume is $V_{\text{cap}} - V_{\text{ball}} = \frac{1}{3}\pi(45h^2 - h^3 - 500)$ cm^3.





83. Take the x-axis to be the axis of the cylindrical hole of radius r. A quarter of the cross-section through y, perpendicular to the y-axis, is the rectangle shown. Using the Pythagorean Theorem twice, we see that the dimensions of this rectangle are $x = \sqrt{R^2 - y^2}$ and $z = \sqrt{r^2 - y^2}$, so $\frac{1}{4}A(y) = xz = \sqrt{r^2 - y^2}\,\sqrt{R^2 - y^2}$, and



$$V = \int_{-r}^{r} A(y)\,dy = \int_{-r}^{r} 4\sqrt{r^2 - y^2}\,\sqrt{R^2 - y^2}\,dy = 8\int_0^r \sqrt{r^2 - y^2}\,\sqrt{R^2 - y^2}\,dy$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

84. The line $y = r$ intersects the semicircle $y = \sqrt{R^2 - x^2}$ when $r = \sqrt{R^2 - x^2} \;\Rightarrow\; r^2 = R^2 - x^2 \;\Rightarrow$ $x^2 = R^2 - r^2 \;\Rightarrow\; x = \pm\sqrt{R^2 - r^2}$. Rotating the shaded region about the x-axis gives us

$$\begin{aligned} V &= \int_{-\sqrt{R^2-r^2}}^{\sqrt{R^2-r^2}} \pi\left[\left(\sqrt{R^2-x^2}\right)^2 - r^2\right] dx = 2\pi \int_0^{\sqrt{R^2-r^2}} (R^2 - x^2 - r^2)\, dx \qquad \text{[by symmetry]} \\ &= 2\pi \int_0^{\sqrt{R^2-r^2}} \left[\left(R^2 - r^2\right) - x^2\right] dx = 2\pi\left[\left(R^2 - r^2\right)x - \tfrac{1}{3}x^3\right]_0^{\sqrt{R^2-r^2}} \\ &= 2\pi\left[\left(R^2 - r^2\right)^{3/2} - \tfrac{1}{3}\left(R^2 - r^2\right)^{3/2}\right] = 2\pi \cdot \tfrac{2}{3}\left(R^2 - r^2\right)^{3/2} = \tfrac{4\pi}{3}\left(R^2 - r^2\right)^{3/2} \end{aligned}$$

Our answer makes sense in limiting cases. As $r \to 0$, $V \to \frac{4}{3}\pi R^3$, which is the volume of the full sphere. As $r \to R$, $V \to 0$, which makes sense because the hole's radius is approaching that of the sphere.





85. (a) The radius of the barrel is the same at each end by symmetry, since the function $y = R - cx^2$ is even. Since the barrel is obtained by rotating the graph of the function y about the x-axis, this radius is equal to the value of y at $x = \frac{1}{2}h$, which is $R - c\left(\frac{1}{2}h\right)^2 = R - d = r$.



(b) The barrel is symmetric about the y-axis, so its volume is twice the volume of that part of the barrel for $x > 0$. Also, the barrel is a volume of rotation, so

$$\begin{aligned} V &= 2\int_0^{h/2} \pi y^2\, dx = 2\pi \int_0^{h/2} \left(R - cx^2\right)^2 dx = 2\pi\left[R^2 x - \tfrac{2}{3}Rcx^3 + \tfrac{1}{5}c^2x^5\right]_0^{h/2} \\ &= 2\pi\left(\tfrac{1}{2}R^2h - \tfrac{1}{12}Rch^3 + \tfrac{1}{160}c^2h^5\right) \end{aligned}$$

Trying to make this look more like the expression we want, we rewrite it as $V = \frac{1}{3}\pi h\left[2R^2 + \left(R^2 - \frac{1}{2}Rch^2 + \frac{3}{80}c^2h^4\right)\right]$. But $R^2 - \frac{1}{2}Rch^2 + \frac{3}{80}c^2h^4 = \left(R - \frac{1}{4}ch^2\right)^2 - \frac{1}{40}c^2h^4 = (R - d)^2 - \frac{2}{5}\left(\frac{1}{4}ch^2\right)^2 = r^2 - \frac{2}{5}d^2$.

Substituting this back into V, we see that $V = \frac{1}{3}\pi h\left(2R^2 + r^2 - \frac{2}{5}d^2\right)$, as required.

86. It suffices to consider the case where $\mathcal{R}$ is bounded by the curves $y = f(x)$ and $y = g(x)$ for $a \le x \le b$, where $g(x) \le f(x)$ for all x in $[a, b]$, since other regions can be decomposed into subregions of this type. We are concerned with the volume obtained when $\mathcal{R}$ is rotated about the line $y = -k$, which is equal to

$$\begin{aligned} V_2 &= \pi \textstyle\int_a^b \left([f(x) + k]^2 - [g(x) + k]^2\right) dx \\ &= \pi \textstyle\int_a^b \left([f(x)]^2 - [g(x)]^2\right) dx + 2\pi k \int_a^b [f(x) - g(x)]\, dx = V_1 + 2\pi kA \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

87. (a) $y = x^3 \quad \Rightarrow \quad x = \sqrt[3]{y}$. $V_1 = \int_1^8 \pi \left(\sqrt[3]{y}\right)^2 \, dy = \int_1^8 \pi y^{2/3} \, dy = \frac{3}{5}\pi \left[y^{5/3}\right]_1^8 = \frac{3}{5}\pi(32-1) = \frac{93}{5}\pi$

(b) If each y is replaced with cy, then $y = 1$ will be mapped to $y = 1c = c$, and $y = 8$ will be mapped to $y = 8c$, each as shown. If each x is replaced with cx, then $y = (cx)^3/c^2 = c^3x^3/c^2 = cx^3$, so y has again been mapped to cy. A dilation with scaling factor c therefore transforms the region R_1 into R_2.

(c) $y = x^3/c^2 \quad \Rightarrow \quad c^2y = x^3 \quad \Rightarrow \quad x = \sqrt[3]{c^2y}$. Then

$$V_2 = \int_c^{8c} \pi\left(\sqrt[3]{c^2y}\right)^2 \, dy = \int_c^{8c} \pi \cdot c^{4/3}y^{2/3} \, dy = \frac{3}{5}\pi \cdot c^{4/3}\left[y^{5/3}\right]_c^{8c} = \frac{3}{5}\pi \cdot c^{4/3}(32c^{5/3} - c^{5/3})$$
$$= \frac{3}{5}\pi \cdot c^{4/3} \cdot 31c^{5/3} = \frac{3}{5}\pi \cdot 31c^3 = \frac{93}{5}\pi c^3 = c^3V_1$$

(d) $V_2 = 5 \text{ L} \quad \Rightarrow \quad \frac{93}{5}\pi c^3 = 5000 \text{ cm}^3 \quad \Rightarrow \quad c^3 = \frac{25{,}000}{93\pi} \quad \Rightarrow \quad c = \sqrt[3]{\frac{25{,}000}{93\pi}} \approx 4.41$

6.3 Volumes by Cylindrical Shells

1.



If we were to use the "washer" method, we would first have to locate the local maximum point (a, b) of $y = x(x-1)^2$ using the methods of Chapter 4. Then we would have to solve the equation $y = x(x-1)^2$ for x in terms of y to obtain the functions $x = g_1(y)$ and $x = g_2(y)$ shown in the first figure. This step would be difficult because it involves the cubic formula. Finally we would find the volume using $V = \pi \int_0^b \left\{[g_1(y)]^2 - [g_2(y)]^2\right\} dy$.

Using shells, we find that a typical approximating shell has radius x, so its circumference is $2\pi x$. Its height is y, that is, $x(x-1)^2$. So the total volume is

$$V = \int_0^1 2\pi x\left[x(x-1)^2\right] dx = 2\pi \int_0^1 \left(x^4 - 2x^3 + x^2\right) dx = 2\pi\left[\frac{x^5}{5} - 2\frac{x^4}{4} + \frac{x^3}{3}\right]_0^1 = \frac{\pi}{15}$$

2.



A typical cylindrical shell has circumference $2\pi x$ and height $\sin(x^2)$. $V = \int_0^{\sqrt{\pi}} 2\pi x \sin(x^2) \, dx$. Let $u = x^2$. Then $du = 2x \, dx$, so $V = \pi \int_0^{\pi} \sin u \, du = \pi[-\cos u]_0^{\pi} = \pi[1 - (-1)] = 2\pi$. For washers, we would first have to locate the local maximum point (a, b) of $y = \sin(x^2)$ using the methods of Chapter 4. Then we would have to solve the equation $y = \sin\left(x^2\right)$ for x in terms of y to obtain the functions $x = g_1(y)$ and $x = g_2(y)$ shown in the second figure. Finally we would find the volume using $V = \pi \int_0^b \left\{[g_1(y)]^2 - [g_2(y)]^2\right\} dy$. Using shells is definitely preferable to using washers.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. (a)





$$V = \int_0^{\sqrt{\pi/2}} 2\pi x \cos(x^2)\,dx$$

(b) $V = \displaystyle\int_0^{\sqrt{\pi/2}} 2\pi x\cos(x^2)\,dx = 2\pi\int_0^{\sqrt{\pi/2}} x\cos(x^2)\,dx = 2\pi\cdot\tfrac{1}{2}\Big[\sin(x^2)\Big]_0^{\sqrt{\pi/2}} = \pi\Big[\sin\Big(\frac{\pi}{2}\Big) - \sin 0\Big]$

$= \pi(1-0) = \pi$

4. (a)





$y = 2 - x \quad\Rightarrow\quad x = 2 - y;$

$y = \sqrt{x} \quad\Rightarrow\quad x = y^2$

$$V = \int_0^1 2\pi y[(2-y) - y^2]\,dy$$

(b) $V = \int_0^1 2\pi y[(2-y)-y^2]\,dy = 2\pi\int_0^1 (2y - y^2 - y^3)\,dy = 2\pi\big[y^2 - \tfrac{1}{3}y^3 - \tfrac{1}{4}y^4\big]_0^1$

$2\pi\big[(1 - \tfrac{1}{3} - \tfrac{1}{4}) - 0\big] = 2\pi\big(\tfrac{5}{12}\big) = \tfrac{5}{6}\pi$

5. $V = \int_1^2 2\pi x \ln x\,dx$



6. $y = x^3 \quad\Rightarrow\quad x = \sqrt[3]{y}$

$V = \int_0^8 2\pi y\sqrt[3]{y}\,dy$



7. $y = \sin^{-1} x \quad\Rightarrow\quad x = \sin y$

$V = \int_0^{\pi/2} 2\pi(3-y)\sin y\,dy$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. $V = \int_0^3 2\pi(7-x)[(4x-x^2)-x]\,dx$



9. The shell has radius x, circumference $2\pi x$, and height $\sqrt{x}$, so

$$V = \int_0^4 2\pi x\sqrt{x}\,dx = \int_0^4 2\pi x^{3/2}\,dx = 2\pi\left[\tfrac{2}{5}x^{5/2}\right]_0^4 = 2\pi\cdot\tfrac{2}{5}(32-0) = \tfrac{128}{5}\pi.$$





10. The shell has radius x, circumference $2\pi x$, and height x^3, so

$$V = \int_1^2 2\pi x\cdot x^3\,dx = 2\pi\int_1^2 x^4\,dx$$
$$= 2\pi\left[\tfrac{1}{5}x^5\right]_1^2 = 2\pi\left(\tfrac{32}{5}-\tfrac{1}{5}\right) = \tfrac{62}{5}\pi$$





11. The shell has radius x, circumference $2\pi x$, and height $1/x$, so

$$V = \int_1^4 2\pi x\left(\frac{1}{x}\right)dx = \int_1^4 2\pi\,dx = 2\pi\left[x\right]_1^4 = 2\pi(4-1) = 6\pi.$$





12. The shell has radius x, circumference $2\pi x$, and height e^{-x^2}, so $V = \int_0^1 2\pi x e^{-x^2}\,dx$.

Let $u = x^2$. Thus, $du = 2x\,dx$, so

$$V = \pi\int_0^1 e^{-u}\,du = \pi\left[-e^{-u}\right]_0^1 = \pi(1-1/e).$$





© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

13. The shell has radius x, circumference $2\pi x$, and height $\sqrt{5+x^2}$, so

$$\begin{aligned} V &= \int_0^2 2\pi x\sqrt{5+x^2}\,dx \quad [\,u = 5+x^2,\, du = 2x\,dx\,] \\ &= \int_5^9 2\pi \cdot \frac{1}{2}u^{1/2}\,du = \pi \cdot \frac{2}{3}\Big[u^{3/2}\Big]_5^9 \\ &= \frac{2}{3}\pi(27 - 5^{3/2}) \end{aligned}$$





14. $4x - x^2 = x \;\Leftrightarrow\; 0 = x^2 - 3x \;\Leftrightarrow\; 0 = x(x-3) \;\Leftrightarrow\; x = 0$ or 3.

The shell has radius x, circumference $2\pi x$, and height $[(4x - x^2) - x]$, so

$$\begin{aligned} V &= \int_0^3 2\pi x[(4x - x^2) - x]\,dx \\ &= 2\pi \int_0^3 (-x^3 + 3x^2)\,dx \\ &= 2\pi\big[-\tfrac{1}{4}x^4 + x^3\big]_0^3 \\ &= 2\pi\big(-\tfrac{81}{4} + 27\big) = 2\pi\big(\tfrac{27}{4}\big) = \tfrac{27}{2}\pi \end{aligned}$$





15. $xy = 1 \;\Rightarrow\; x = \frac{1}{y}$. The shell has radius y, circumference $2\pi y$, and height $1/y$, so

$$\begin{aligned} V &= \int_1^3 2\pi y\left(\frac{1}{y}\right) dy \\ &= 2\pi \int_1^3 dy = 2\pi\big[y\big]_1^3 \\ &= 2\pi(3-1) = 4\pi \end{aligned}$$





16. $y = \sqrt{x} \;\Rightarrow\; x = y^2$. The shell has radius y, circumference $2\pi y$, and height y^2, so

$$\begin{aligned} V &= \int_0^2 2\pi y(y^2)\,dy = 2\pi \int_0^2 y^3\,dy \\ &= 2\pi\big[\tfrac{1}{4}y^4\big]_0^2 \\ &= 2\pi(4) = 8\pi \end{aligned}$$





© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. $y = x^{3/2} \quad \Rightarrow \quad x = y^{2/3}$. The shell has radius y, circumference $2\pi y$, and height $y^{2/3}$, so

$$V = \int_0^8 2\pi y(y^{2/3})\,dy = 2\pi \int_0^8 y^{5/3}\,dy$$
$$= 2\pi\left[\tfrac{3}{8}y^{8/3}\right]_0^8$$
$$= 2\pi \cdot \tfrac{3}{8} \cdot 256 = 192\pi$$



18. The shell has radius y, circumference $2\pi y$, and height $-3y^2 + 12y - 9$, so

$$V = \int_1^3 2\pi y(-3y^2 + 12y - 9)\,dy$$
$$= 2\pi \int_1^3 (-3y^3 + 12y^2 - 9y)\,dy$$
$$= -6\pi \int_1^3 (y^3 - 4y^2 + 3y)\,dy$$
$$= -6\pi\left[\tfrac{1}{4}y^4 - \tfrac{4}{3}y^3 + \tfrac{3}{2}y^2\right]_1^3$$
$$= -6\pi\left[\left(\tfrac{81}{4} - 36 + \tfrac{27}{2}\right) - \left(\tfrac{1}{4} - \tfrac{4}{3} + \tfrac{3}{2}\right)\right]$$
$$= -6\pi\left(-\tfrac{8}{3}\right) = 16\pi$$



19. The shell has radius y, circumference $2\pi y$, and height

$$2 - \left[1 + (y-2)^2\right] = 1 - (y-2)^2 = 1 - \left(y^2 - 4y + 4\right)$$
$$= -y^2 + 4y - 3, \text{ so}$$

$$V = \int_1^3 2\pi y(-y^2 + 4y - 3)\,dy$$
$$= 2\pi \int_1^3 (-y^3 + 4y^2 - 3y)\,dy$$
$$= 2\pi\left[-\tfrac{1}{4}y^4 + \tfrac{4}{3}y^3 - \tfrac{3}{2}y^2\right]_1^3$$
$$= 2\pi\left[\left(-\tfrac{81}{4} + 36 - \tfrac{27}{2}\right) - \left(-\tfrac{1}{4} + \tfrac{4}{3} - \tfrac{3}{2}\right)\right]$$
$$= 2\pi\left(\tfrac{8}{3}\right) = \tfrac{16}{3}\pi$$



20. The curves intersect when $4 - y = y^2 - 4y + 4 \quad \Leftrightarrow$

$0 = y^2 - 3y \quad \Leftrightarrow \quad 0 = y(y-3) \quad \Leftrightarrow \quad y = 0$ or 3.

The shell has radius y, circumference $2\pi y$, and height

$(4 - y) - (y^2 - 4y + 4) = -y^2 + 3y$, so

$$V = \int_0^3 2\pi y(-y^2 + 3y)\,dy = 2\pi \int_0^3 (3y^2 - y^3)\,dy$$
$$= 2\pi\left[y^3 - \tfrac{1}{4}y^4\right]_0^3 = 2\pi\left(27 - \tfrac{81}{4}\right) = 2\pi\left(\tfrac{27}{4}\right) = \tfrac{27\pi}{2}$$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

21. $x^2 = 8\sqrt{x} \;\Rightarrow\; x^4 = 64x \;\Rightarrow\; x^4 - 64x = 0 \;\Rightarrow$
$x(x^3 - 64) = 0 \;\Rightarrow\; x = 0$ or $x = 4$



(a) By cylindrical shells:



$$\begin{aligned} V &= \int_0^4 2\pi x \left(8\sqrt{x} - x^2\right) dx \\ &= \int_0^4 2\pi(8x^{3/2} - x^3)\, dx \\ &= 2\pi\left[\tfrac{16}{5}x^{5/2} - \tfrac{1}{4}x^4\right]_0^4 \\ &= 2\pi\left[\left(\tfrac{512}{5} - 64\right) - 0\right] = \tfrac{384}{5}\pi \end{aligned}$$

(b) By washers:



$$\begin{aligned} V &= \int_0^{16} \pi\left[\left(\sqrt{y}\right)^2 - \left(\tfrac{1}{64}y^2\right)^2\right] dy \\ &= \int_0^{16} \pi\left(y - \tfrac{1}{4096}y^4\right) dy = \pi\left[\tfrac{1}{2}y^2 - \tfrac{1}{20{,}480}y^5\right]_0^{16} \\ &= \pi\left(128 - \tfrac{256}{5}\right) = \tfrac{384}{5}\pi \end{aligned}$$

22. $x^3 = 4x^2 \;\Rightarrow\; x^3 - 4x^2 = 0 \;\Rightarrow\; x^2(x - 4) = 0 \;\Rightarrow$
$x = 0$ or $x = 4$



(a) By washers:



$$\begin{aligned} V &= \int_0^4 \pi[(4x^2)^2 - (x^3)^2]\, dx = \int_0^4 \pi(16x^4 - x^6)\, dx \\ &= \pi\left[\frac{16}{5}x^5 - \frac{1}{7}x^7\right]_0^4 = \pi\left(\frac{16{,}384}{5} - \frac{16{,}384}{7}\right) \\ &= \frac{32{,}768}{35}\pi \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) By cylindrical shells:

$$V = \int_0^{64} 2\pi y\left(\sqrt[3]{y} - \tfrac{1}{2}\sqrt{y}\right) dy = \int_0^{64} 2\pi\left(y^{4/3} - \tfrac{1}{2}y^{3/2}\right) dy$$
$$= 2\pi\left[\frac{3}{7}y^{7/3} - \frac{1}{2}\cdot\frac{2}{5}y^{5/2}\right]_0^{64}$$
$$= 2\pi\left(\frac{49{,}152}{7} - \frac{32{,}768}{5}\right) = \frac{32{,}768}{35}\pi$$



23. (a) The shell has radius $x - (-2) = x + 2$, circumference $2\pi(x+2)$, and height $4x - x^2$.



(b) $V = \int_0^4 2\pi(x+2)(4x - x^2)\,dx$

(c) $V = \int_0^4 2\pi(x+2)(4x - x^2)\,dx$
$$= \int_0^4 2\pi(2x^2 - x^3 + 8x)\,dx = 2\pi\left[\frac{2}{3}x^3 - \frac{1}{4}x^4 + 4x^2\right]_0^4$$
$$= 2\pi\left[\left(\frac{128}{3} - 64 + 64\right) - 0\right] = \frac{256}{3}\pi$$

24. (a) $y = \sqrt{x} \;\Rightarrow\; x = y^2$; $y = x^3 \;\Rightarrow\; x = \sqrt[3]{y}$.
The shell has radius $y - (-1) = y + 1$, circumference $2\pi(y+1)$, and height $\sqrt[3]{y} - y^2$.



(b) $V = \int_0^1 2\pi(y+1)\left(\sqrt[3]{y} - y^2\right) dy$

(c) $V = \int_0^1 2\pi(y+1)\left(\sqrt[3]{y} - y^2\right) dy$
$$= \int_0^1 2\pi(y^{4/3} - y^3 + y^{1/3} - y^2)\,dy$$
$$= 2\pi\left[\tfrac{3}{7}y^{7/3} - \tfrac{1}{4}y^4 + \tfrac{3}{4}y^{4/3} - \tfrac{1}{3}y^3\right]_0^1 = 2\pi\left[\left(\tfrac{3}{7} - \tfrac{1}{4} + \tfrac{3}{4} - \tfrac{1}{3}\right) - 0\right]$$
$$= \tfrac{25}{21}\pi$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

25. The shell has radius $3-x$, circumference $2\pi(3-x)$, and height $8-x^3$.

$$\begin{aligned} V &= \int_0^2 2\pi(3-x)(8-x^3)\,dx \\ &= 2\pi\int_0^2 (x^4-3x^3-8x+24)\,dx \\ &= 2\pi\left[\tfrac{1}{5}x^5-\tfrac{3}{4}x^4-4x^2+24x\right]_0^2 \\ &= 2\pi\left(\tfrac{32}{5}-12-16+48\right) = 2\pi\left(\tfrac{132}{5}\right) = \tfrac{264\pi}{5} \end{aligned}$$



26. The shell has radius $x-(-1)=x+1$, circumference $2\pi(x+1)$, and height $4-2x$.

$$\begin{aligned} V &= \int_0^2 2\pi(x+1)(4-2x)\,dx \\ &= 4\pi\int_0^2 (x+1)(2-x)\,dx \\ &= 4\pi\int_0^2 (-x^2+x+2)\,dx \\ &= 4\pi\left[-\tfrac{1}{3}x^3+\tfrac{1}{2}x^2+2x\right]_0^2 \\ &= 4\pi\left(-\tfrac{8}{3}+2+4\right) = 4\pi\left(\tfrac{10}{3}\right) = \tfrac{40\pi}{3} \end{aligned}$$



27. The shell has radius $x-1$, circumference $2\pi(x-1)$, and height $(4x-x^2)-3=-x^2+4x-3$.

$$\begin{aligned} V &= \int_1^3 2\pi(x-1)(-x^2+4x-3)\,dx \\ &= 2\pi\int_1^3 (-x^3+5x^2-7x+3)\,dx \\ &= 2\pi\left[-\tfrac{1}{4}x^4+\tfrac{5}{3}x^3-\tfrac{7}{2}x^2+3x\right]_1^3 \\ &= 2\pi\left[\left(-\tfrac{81}{4}+45-\tfrac{63}{2}+9\right)-\left(-\tfrac{1}{4}+\tfrac{5}{3}-\tfrac{7}{2}+3\right)\right] \\ &= 2\pi\left(\tfrac{4}{3}\right) = \tfrac{8}{3}\pi \end{aligned}$$



28. The shell has radius $5-x$, circumference $2\pi(5-x)$, and height $\sqrt{x}-\frac{1}{2}x$.

$$\begin{aligned} V &= \int_0^4 2\pi(5-x)\left(\sqrt{x}-\tfrac{1}{2}x\right)\,dx = 2\pi\int_0^4 (5x^{1/2}-\tfrac{5}{2}x-x^{3/2}+\tfrac{1}{2}x^2)\,dx \\ &= 2\pi\left[\tfrac{10}{3}x^{3/2}-\tfrac{5}{4}x^2-\tfrac{2}{5}x^{5/2}+\tfrac{1}{6}x^3\right]_0^4 = 2\pi\left(\tfrac{80}{3}-20-\tfrac{64}{5}+\tfrac{32}{3}\right) \\ &= 2\pi\left(\tfrac{68}{15}\right) = \tfrac{136\pi}{15} \end{aligned}$$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

29. The shell has radius $2-y$, circumference $2\pi(2-y)$, and height $2-2y^2$.

$$\begin{aligned} V &= \int_0^1 2\pi(2-y)(2-2y^2)\,dy \\ &= 4\pi\int_0^1 (2-y)(1-y^2)\,dy \\ &= 4\pi\int_0^1 (y^3-2y^2-y+2)\,dy \\ &= 4\pi\left[\tfrac{1}{4}y^4-\tfrac{2}{3}y^3-\tfrac{1}{2}y^2+2y\right]_0^1 \\ &= 4\pi\left(\tfrac{1}{4}-\tfrac{2}{3}-\tfrac{1}{2}+2\right) \\ &= 4\pi\left(\tfrac{13}{12}\right) = \tfrac{13\pi}{3} \end{aligned}$$





30. The shell has radius $y-(-2)=y+2$, circumference $2\pi(y+2)$, and height $(y^2+1)-2y^2=1-y^2$.

$$\begin{aligned} V &= \int_{-1}^1 2\pi(y+2)(1-y^2)\,dy \\ &= 2\pi\int_{-1}^1 (-y^3-2y^2+y+2)\,dy \\ &= 4\pi\int_0^1 (-2y^2+2)\,dy \qquad \text{[by Theorem 5.5.7]} \\ &= 8\pi\int_0^1 (1-y^2)\,dy = 8\pi\left[y-\tfrac{1}{3}y^3\right]_0^1 \\ &= 8\pi\left(1-\tfrac{1}{3}\right) = 8\pi\left(\tfrac{2}{3}\right) = \tfrac{16\pi}{3} \end{aligned}$$





31. (a) $V = 2\pi\displaystyle\int_0^2 x(xe^{-x})\,dx = 2\pi\int_0^2 x^2e^{-x}\,dx$

(b) $V \approx 4.06300$



32. (a) $V = 2\pi\displaystyle\int_0^{\pi/4}\left(\frac{\pi}{2}-x\right)\tan x\,dx$

(b) $V \approx 2.25323$



33. (a) $V = 2\pi\displaystyle\int_{-\pi/2}^{\pi/2}(\pi-x)[\cos^4x-(-\cos^4x)]\,dx$

$$= 4\pi\int_{-\pi/2}^{\pi/2}(\pi-x)\,\cos^4x\,dx$$

[or $8\pi^2\int_0^{\pi/2}\cos^4x\,dx$ using Theorem 5.5.7]

(b) $V \approx 46.50942$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

34. (a) $x = \dfrac{2x}{1+x^3} \quad \Rightarrow \quad x + x^4 = 2x \quad \Rightarrow \quad x^4 - x = 0 \quad \Rightarrow$

$x(x^3 - 1) = 0 \quad \Rightarrow \quad x(x-1)(x^2 + x + 1) = 0 \quad \Rightarrow \quad x = 0 \text{ or } 1$

$V = 2\pi \displaystyle\int_0^1 [x - (-1)] \left(\frac{2x}{1+x^3} - x \right) dx$



(b) $V \approx 2.36164$

35. (a) $V = \int_0^{\pi} 2\pi(4 - y)\sqrt{\sin y}\, dy$ (b) $V \approx 36.57476$



36. (a) $V = \int_{-3}^{3} 2\pi(5 - y)\left(4 - \sqrt{y^2 + 7}\right) dy$ (b) $V \approx 163.02712$



37. $V = \int_0^1 2\pi x\sqrt{1 + x^3}\, dx$. Let $f(x) = x\sqrt{1 + x^3}$.

Then the Midpoint Rule with $n = 5$ gives

$\int_0^1 f(x)\, dx \approx \frac{1-0}{5}[f(0.1) + f(0.3) + f(0.5) + f(0.7) + f(0.9)]$

$\approx 0.2(2.9290)$

Multiplying by 2π gives $V \approx 3.68$.



38. $V = \int_0^{10} 2\pi x f(x)\, dx$. Let $g(x) = xf(x)$, where the values of f are obtained from the graph.

Using the Midpoint Rule with $n = 5$ gives

$$\int_0^{10} g(x)\, dx \approx \tfrac{10-0}{5}[g(1) + g(3) + g(5) + g(7) + g(9)] = 2[1f(1) + 3f(3) + 5f(5) + 7f(7) + 9f(9)]$$

$$= 2[1(4-2) + 3(5-1) + 5(4-1) + 7(4-2) + 9(4-2)]$$

$$= 2(2 + 12 + 15 + 14 + 18) = 2(61) = 122$$

Multiplying by 2π gives $V \approx 244\pi \approx 766.5$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

39. $\int_0^3 2\pi x^5\,dx = 2\pi\int_0^3 x(x^4)\,dx$. By the method of cylindrical shells, this integral represents the volume of the solid obtained by rotating the region $0 \le y \le x^4$, $0 \le x \le 3$ about the y-axis ($x = 0$).

40. $\int_1^3 2\pi y \ln y\,dy$. By the method of cylindrical shells, this integral represents the volume of the solid obtained by rotating the region $0 \le x \le \ln y$, $1 \le y \le 3$ about the x-axis.

41. $2\pi\displaystyle\int_1^4 \frac{y+2}{y^2}\,dy = 2\pi\int_1^4 (y+2)\left(\frac{1}{y^2}\right)dy$. By the method of cylindrical shells, this integral represents the volume of the solid obtained by rotating the region $0 \le x \le 1/y^2$, $1 \le y \le 4$ about the line $y = -2$.

42. $\int_0^1 2\pi(2-x)(3^x - 2^x)\,dx$. By the method of cylindrical shells, this integral represents the volume of the solid obtained by rotating the region $2^x \le y \le 3^x$, $0 \le x \le 1$ about the line $x = 2$.

43. From the graph, the curves intersect at $x = 0$ and $x = a \approx 2.175$, with $\dfrac{x}{x^2+1} > x^2 - 2x$ on the interval $(0, a)$. So the volume of the solid obtained by rotating the region about the y-axis is

$$V = 2\pi\int_0^a x\left[\frac{x}{x^2+1} - (x^2 - 2x)\right]dx \approx 14.450$$



44. From the graph, the curves intersect at $x = a \approx 0.906$ and $x = b \approx 2.715$, with $e^{\sin x} > x^2 - 4x + 5$ on the interval (a, b). So the volume of the solid obtained by rotating the region about the y-axis is

$$V = 2\pi\int_a^b x\left[e^{\sin x} - (x^2 - 4x + 5)\right]dx \approx 21.253$$



45. $V = 2\pi\displaystyle\int_0^{\pi/2}\left[\left(\tfrac{\pi}{2} - x\right)\left(\sin^2 x - \sin^4 x\right)\right]dx$

$\overset{\text{CAS}}{=} \frac{1}{32}\pi^3$



46. $V = 2\pi\int_0^{\pi}\left\{[x - (-1)](x^3 \sin x)\right\}dx$

$\overset{\text{CAS}}{=} 2\pi(\pi^4 + \pi^3 - 12\pi^2 - 6\pi + 48)$

$= 2\pi^5 + 2\pi^4 - 24\pi^3 - 12\pi^2 + 96\pi$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

47. (a) Use shells. Each shell has radius x, circumference $2\pi x$, and height $\dfrac{1}{1+x^2}-\dfrac{x}{2}$.

$$V=\int_0^1 2\pi x\left(\frac{1}{1+x^2}-\frac{x}{2}\right)dx.$$

(b) $V=\displaystyle\int_0^1 2\pi x\left(\frac{1}{1+x^2}-\frac{x}{2}\right)dx=\int_0^1 2\pi\left(\frac{x}{1+x^2}-\frac{1}{2}x^2\right)dx=2\pi\left[\frac{1}{2}\ln\left|1+x^2\right|-\frac{1}{6}x^3\right]_0^1$

$=2\pi\left[\left(\frac{1}{2}\ln 2-\frac{1}{6}\right)-0\right]=\pi\left(\ln 2-\frac{1}{3}\right)$

48. (a) Use washers. $2-x^2=x^2 \;\Rightarrow\; 2x^2=2 \;\Rightarrow\; x=1 \;\; [x>0]$

$V=\int_0^1 \pi[(2-x^2)^2-(x^2)^2]\,dx$

(b) $V=\int_0^1 \pi[(2-x^2)^2-(x^2)^2]\,dx=\int_0^1 \pi(4-4x^2)\,dx=\pi\left[4x-4\cdot\frac{1}{3}x^3\right]_0^1=4\pi\left[\left(1-\frac{1}{3}\right)-0\right]=\frac{8}{3}\pi$

49. (a) Use disks. $V=\int_0^\pi \pi\left(\sqrt{\sin x}\right)^2\,dx$

(b) $V=\int_0^\pi \pi\left(\sqrt{\sin x}\right)^2\,dx=\int_0^\pi \pi\sin x\,dx=\pi\Big[-\cos x\Big]_0^\pi=\pi[-\cos\pi-(-\cos 0)]=\pi(1+1)=2\pi$

50. (a) Use shells. Each shell has radius x, circumference $2\pi x$, and height $[(4x-x^2)-x]$.

$4x-x^2=x \;\Rightarrow\; x^2-3x=0 \;\Rightarrow\; x(x-3)=0 \;\Rightarrow\; x=0$ or $x=3$

$V=\int_0^3 2\pi x[(4x-x^2)-x]\,dx$

(b) $V=\int_0^3 2\pi x[(4x-x^2)-x]\,dx=\int_0^3 2\pi(3x^2-x^3)\,dx=2\pi\left[x^3-\frac{1}{4}x^4\right]_0^3=2\pi\left[\left(27-\frac{81}{4}\right)-0\right]=\frac{27}{2}\pi$

51. (a) Use shells. Each shell has radius $x-(-2)=x+2$, circumference $2\pi(x+2)$, and height x^2-x^3.

$V=\int_0^{1/2} 2\pi(x+2)(x^2-x^3)\,dx$

(b) $V=\displaystyle\int_0^{1/2} 2\pi(x+2)(x^2-x^3)\,dx=\int_0^{1/2} 2\pi(-x^3-x^4+2x^2)\,dx=2\pi\left[-\frac{1}{4}x^4-\frac{1}{5}x^5+\frac{2}{3}x^3\right]_0^{1/2}$

$=2\pi\left[\left(-\frac{1}{64}-\frac{1}{160}+\frac{1}{12}\right)-0\right]=\frac{59}{480}\pi$

52. (a) Use shells. Each shell has radius $3-y$, circumference $2\pi(3-y)$, and height $[(3y-y^2)-2]$.

$3y-y^2=2 \;\Rightarrow\; y^2-3y+2=0 \;\Rightarrow\; (y-1)(y-2)=0 \;\Rightarrow\; y=1$ or $y=2$

$V=\int_1^2 2\pi(3-y)[(3y-y^2)-2]\,dy$

(b) $V=\displaystyle\int_1^2 2\pi(3-y)[(3y-y^2)-2]\,dy=\int_1^2 2\pi(y^3-6y^2+11y-6)\,dy$

$=2\pi\left[\frac{1}{4}y^4-2y^3+\frac{11}{2}y^2-6y\right]_1^2=2\pi\left[(4-16+22-12)-\left(\frac{1}{4}-2+\frac{11}{2}-6\right)\right]=\frac{1}{2}\pi$

53. Use shells:

$V=\int_2^4 2\pi x(-x^2+6x-8)\,dx=2\pi\int_2^4(-x^3+6x^2-8x)\,dx$

$=2\pi\left[-\frac{1}{4}x^4+2x^3-4x^2\right]_2^4$

$=2\pi[(-64+128-64)-(-4+16-16)]$

$=2\pi(4)=8\pi$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

54. Use disks:

$$\begin{aligned} V &= \int_2^4 \pi(-x^2+6x-8)^2\,dx \\ &= \pi\int_2^4 (x^4-12x^3+52x^2-96x+64)\,dx \\ &= \pi\left[\tfrac{1}{5}x^5-3x^4+\tfrac{52}{3}x^3-48x^2+64x\right]_2^4 \\ &= \pi\left(\tfrac{512}{15}-\tfrac{496}{15}\right)=\tfrac{16}{15}\pi \end{aligned}$$



55. Use washers: $y^2-x^2=1 \quad\Rightarrow\quad y=\pm\sqrt{x^2\pm 1}$

$$\begin{aligned} V &= \int_{-\sqrt{3}}^{\sqrt{3}} \pi\left[(2-0)^2-\left(\sqrt{x^2+1}-0\right)^2\right]dx \\ &= 2\pi\int_0^{\sqrt{3}} [4-(x^2+1)]\,dx \qquad \text{[by symmetry]} \\ &= 2\pi\int_0^{\sqrt{3}} (3-x^2)\,dx = 2\pi\left[3x-\tfrac{1}{3}x^3\right]_0^{\sqrt{3}} \\ &= 2\pi\left(3\sqrt{3}-\sqrt{3}\right)=4\sqrt{3}\,\pi \end{aligned}$$



56. Use disks: $y^2-x^2=1 \quad\Rightarrow\quad x=\pm\sqrt{y^2-1}$

$$\begin{aligned} V &= \pi\int_1^2 \left(\sqrt{y^2-1}\right)^2 dy = \pi\int_1^2 (y^2-1)\,dy \\ &= \pi\left[\tfrac{1}{3}y^3-y\right]_1^2 = \pi\left[\left(\tfrac{8}{3}-2\right)-\left(\tfrac{1}{3}-1\right)\right]=\tfrac{4}{3}\pi \end{aligned}$$



57. Use disks: $x^2+(y-1)^2=1 \quad\Leftrightarrow\quad x=\pm\sqrt{1-(y-1)^2}$

$$\begin{aligned} V &= \pi\int_0^2 \left[\sqrt{1-(y-1)^2}\right]^2 dy = \pi\int_0^2 (2y-y^2)\,dy \\ &= \pi\left[y^2-\tfrac{1}{3}y^3\right]_0^2 = \pi\left(4-\tfrac{8}{3}\right)=\tfrac{4}{3}\pi \end{aligned}$$



58. Use shells:

$$\begin{aligned} V &= \int_1^5 2\pi(y-1)[4-(y-3)^2]\,dy \\ &= 2\pi\int_1^5 (y-1)(-y^2+6y-5)\,dy \\ &= 2\pi\int_1^5 (-y^3+7y^2-11y+5)\,dy \\ &= 2\pi\left[-\tfrac{1}{4}y^4+\tfrac{7}{3}y^3-\tfrac{11}{2}y^2+5y\right]_1^5 \\ &= 2\pi\left(\tfrac{275}{12}-\tfrac{19}{12}\right)=\tfrac{128}{3}\pi \end{aligned}$$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

59. $y+1=(y-1)^2 \quad \Leftrightarrow \quad y+1=y^2-2y+1 \quad \Leftrightarrow \quad 0=y^2-3y \quad \Leftrightarrow$
$0=y(y-3) \quad \Leftrightarrow \quad y=0$ or 3.



Use washers:

$$V=\pi\int_0^3 \left\{[(y+1)-(-1)]^2-[(y-1)^2-(-1)]^2\right\}dy$$
$$=\pi\int_0^3 [(y+2)^2-(y^2-2y+2)^2]\,dy$$
$$=\pi\int_0^3 [(y^2+4y+4)-(y^4-4y^3+8y^2-8y+4)]\,dy=\pi\int_0^3 (-y^4+4y^3-7y^2+12y)\,dy$$
$$=\pi\left[-\tfrac{1}{5}y^5+y^4-\tfrac{7}{3}y^3+6y^2\right]_0^3=\pi\left(-\tfrac{243}{5}+81-63+54\right)=\tfrac{117}{5}\pi$$

60. Use cylindrical shells to find the volume V.



$$V=\int_0^1 2\pi(a-x)(2x)\,dx=4\pi\int_0^1 (ax-x^2)\,dx$$
$$=4\pi\left[\tfrac{1}{2}ax^2-\tfrac{1}{3}x^3\right]_0^1=4\pi\left(\tfrac{1}{2}a-\tfrac{1}{3}\right)$$

Now solve for a in terms of V:

$$V=4\pi\left(\tfrac{1}{2}a-\tfrac{1}{3}\right) \quad \Leftrightarrow \quad \frac{V}{4\pi}=\frac{1}{2}a-\frac{1}{3} \quad \Leftrightarrow \quad \frac{1}{2}a=\frac{V}{4\pi}+\frac{1}{3} \quad \Leftrightarrow$$
$$a=\frac{V}{2\pi}+\frac{2}{3}$$

61. Use shells:



$$V=2\int_0^r 2\pi x\sqrt{r^2-x^2}\,dx=-2\pi\int_0^r (r^2-x^2)^{1/2}(-2x)\,dx$$
$$=\left[-2\pi\cdot\tfrac{2}{3}(r^2-x^2)^{3/2}\right]_0^r=-\tfrac{4}{3}\pi(0-r^3)=\tfrac{4}{3}\pi r^3$$

62. $V=\int_{R-r}^{R+r} 2\pi x\cdot 2\sqrt{r^2-(x-R)^2}\,dx$



$$=\int_{-r}^{r} 4\pi(u+R)\sqrt{r^2-u^2}\,du \qquad [\text{let } u=x-R]$$
$$=4\pi R\int_{-r}^{r}\sqrt{r^2-u^2}\,du+4\pi\int_{-r}^{r} u\sqrt{r^2-u^2}\,du$$

The first integral is the area of a semicircle of radius r, that is, $\frac{1}{2}\pi r^2$, and the second is zero since the integrand is an odd function. Thus,

$$V=4\pi R\left(\tfrac{1}{2}\pi r^2\right)+4\pi\cdot 0=2\pi^2 Rr^2.$$

63. $V=2\pi\int_0^r x\left(-\frac{h}{r}x+h\right)dx=2\pi h\int_0^r\left(-\frac{x^2}{r}+x\right)dx$



$$=2\pi h\left[-\frac{x^3}{3r}+\frac{x^2}{2}\right]_0^r=2\pi h\,\frac{r^2}{6}=\frac{\pi r^2 h}{3}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

64. By symmetry, the volume of a napkin ring obtained by drilling a hole of radius r through a sphere with radius R is twice the volume obtained by rotating the area above the x-axis and below the curve $y = \sqrt{R^2 - x^2}$ (the equation of the top half of the cross-section of the sphere), between $x = r$ and $x = R$, about the y-axis. This volume is equal to



$$2\int_{\text{inner radius}}^{\text{outer radius}} 2\pi rh\,dx = 2\cdot 2\pi\int_r^R x\sqrt{R^2-x^2}\,dx = 4\pi\left[-\tfrac{1}{3}\left(R^2-x^2\right)^{3/2}\right]_r^R = \tfrac{4}{3}\pi(R^2-r^2)^{3/2}$$

But by the Pythagorean Theorem, $R^2 - r^2 = \left(\frac{1}{2}h\right)^2$, so the volume of the napkin ring is $\frac{4}{3}\pi\left(\frac{1}{2}h\right)^3 = \frac{1}{6}\pi h^3$, which is independent of both R and r; that is, the amount of wood in a napkin ring of height h is the same regardless of the size of the sphere used. Note that most of this calculation has been done already, but with more difficulty, in Exercise 6.2.84.

Another solution: The height of the missing cap is the radius of the sphere minus half the height of the cut-out cylinder, that is, $R - \frac{1}{2}h$. Using Exercise 6.2.61,

$$V_{\text{napkin ring}} = V_{\text{sphere}} - V_{\text{cylinder}} - 2V_{\text{cap}} = \tfrac{4}{3}\pi R^3 - \pi r^2 h - 2\cdot\tfrac{\pi}{3}\left(R-\tfrac{1}{2}h\right)^2\left[3R-\left(R-\tfrac{1}{2}h\right)\right] = \tfrac{1}{6}\pi h^3$$

6.4 Work

1. The force exerted by the weight lifter is $F = mg = (200\text{ kg})(9.8\text{ m/s}^2) = 1960\text{ N}$. The work done by the weight lifter in lifting the weight from 1.5 m to 2.0 m above the ground is then

$$W = Fd = (1960\text{ N})(2.0\text{ m} - 1.5\text{ m}) = (1960\text{ N})(0.5\text{ m}) = 980\text{ N-m} = 980\text{ J}$$

2. $W = Fd = (1100\text{ lb})(35\text{ ft}) = 38{,}500\text{ ft-lb}$.

3. $W = \int_a^b f(x)\,dx = \int_1^{10} 5x^{-2}\,dx = 5\left[-x^{-1}\right]_1^{10} = 5\left(-\frac{1}{10}+1\right) = 4.5\text{ ft-lb}$

4. $W = \displaystyle\int_a^b f(x)\,dx = \int_4^{16} 4\sqrt{x}\,dx = 4\left[\tfrac{2}{3}x^{3/2}\right]_4^{16} = \tfrac{8}{3}(64-8) = \tfrac{448}{3}\text{ N-m} = \tfrac{448}{3}\text{ J}$.

5. The force function is given by $F(x)$ (in newtons) and the work (in joules) is the area under the curve, given by $\int_0^8 F(x)\,dx = \int_0^4 F(x)\,dx + \int_4^8 F(x)\,dx = \frac{1}{2}(4)(30) + (4)(30) = 180\text{ J}$.

6. $W = \int_4^{20} f(x)\,dx \approx M_4 = \Delta x[f(6) + f(10) + f(14) + f(18)] = \frac{20-4}{4}[5.8 + 8.8 + 8.2 + 5.2] = 4(28) = 112\text{ J}$

7. According to Hooke's Law, the force required to maintain a spring stretched x units beyond its natural length (or compressed x units less than its natural length) is proportional to x, that is, $f(x) = kx$. Here, the amount stretched is 4 in. $= \frac{1}{3}$ ft and the force is 10 lb. Thus, $10 = k\left(\frac{1}{3}\right) \quad \Rightarrow \quad k = 30$ lb/ft, and $f(x) = 30x$. The work done in stretching the spring from its natural length to 6 in. $= \frac{1}{2}$ ft beyond its natural length is $W = \int_0^{1/2} 30x\,dx = \left[15x^2\right]_0^{1/2} = \frac{15}{4}$ ft-lb.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

8. According to Hooke's Law, the force required to maintain a spring stretched x units beyond its natural length (or compressed x units less than its natural length) is proportional to x, that is, $f(x) = kx$. Here, the amount compressed is $40 - 30 = 10$ cm $= 0.1$ m and the force is 60 N. Thus, $60 = k(0.1) \quad \Rightarrow \quad k = 600$ N/m, and $f(x) = 600x$. The work required to compress the spring 0.1 m is $W = \int_0^{0.1} 600x\,dx = \left[300x^2\right]_0^{0.1} = 300(0.01) = 3$ N-m (or J). The work required to compress the spring $40 - 25 = 15$ cm $= 0.15$ m is $W = \int_0^{0.15} 600x\,dx = \left[300x^2\right]_0^{0.15} = 300(0.0225) = 6.75$ J.

9. (a) If $\int_0^{0.12} kx\,dx = 2$ J, then $2 = \left[\frac{1}{2}kx^2\right]_0^{0.12} = \frac{1}{2}k(0.0144) = 0.0072k$ and $k = \frac{2}{0.0072} = \frac{2500}{9} \approx 277.78$ N/m.

Thus, the work needed to stretch the spring from 35 cm to 40 cm is

$\int_{0.05}^{0.10} \frac{2500}{9}x\,dx = \left[\frac{1250}{9}x^2\right]_{1/20}^{1/10} = \frac{1250}{9}\left(\frac{1}{100} - \frac{1}{400}\right) = \frac{25}{24} \approx 1.04$ J.

(b) $f(x) = kx$, so $30 = \frac{2500}{9}x$ and $x = \frac{270}{2500}$ m $= 10.8$ cm

10. If $12 = \int_0^1 kx\,dx = \left[\frac{1}{2}kx^2\right]_0^1 = \frac{1}{2}k$, then $k = 24$ lb/ft and the work required is

$\int_0^{3/4} 24x\,dx = \left[12x^2\right]_0^{3/4} = 12 \cdot \frac{9}{16} = \frac{27}{4} = 6.75$ ft-lb.

11. The distance from 20 cm to 30 cm is 0.1 m, so with $f(x) = kx$, we get $W_1 = \int_0^{0.1} kx\,dx = k\left[\frac{1}{2}x^2\right]_0^{0.1} = \frac{1}{200}k$.

Now $W_2 = \int_{0.1}^{0.2} kx\,dx = k\left[\frac{1}{2}x^2\right]_{0.1}^{0.2} = k\left(\frac{4}{200} - \frac{1}{200}\right) = \frac{3}{200}k$. Thus, $W_2 = 3W_1$.

12. Let L be the natural length of the spring in meters. Then

$6 = \int_{0.10-L}^{0.12-L} kx\,dx = \left[\frac{1}{2}kx^2\right]_{0.10-L}^{0.12-L} = \frac{1}{2}k\left[(0.12 - L)^2 - (0.10 - L)^2\right]$ and

$10 = \int_{0.12-L}^{0.14-L} kx\,dx = \left[\frac{1}{2}kx^2\right]_{0.12-L}^{0.14-L} = \frac{1}{2}k\left[(0.14 - L)^2 - (0.12 - L)^2\right]$.

Simplifying gives us $12 = k(0.0044 - 0.04L)$ and $20 = k(0.0052 - 0.04L)$. Subtracting the first equation from the second gives $8 = 0.0008k$, so $k = 10{,}000$. Now the second equation becomes $20 = 52 - 400L$, so $L = \frac{32}{400}$ m $= 8$ cm.

In Exercises 13–22, n is the number of subintervals of length Δx, and x_i^* is a sample point in the ith subinterval $[x_{i-1}, x_i]$.

13. (a) The portion of the rope from x ft to $(x + \Delta x)$ ft below the top of the building weighs $\frac{1}{2}\,\Delta x$ lb and must be lifted x_i^* ft, so its contribution to the total work is $\frac{1}{2}x_i^*\,\Delta x$ ft-lb. The total work is

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} \frac{1}{2}x_i^*\,\Delta x = \int_0^{50} \frac{1}{2}x\,dx = \left[\frac{1}{4}x^2\right]_0^{50} = \frac{2500}{4} = 625 \text{ ft-lb}$$

Notice that the exact height of the building does not matter (as long as it is more than 50 ft).

(b) When half the rope is pulled to the top of the building, the work to lift the top half of the rope is $W_1 = \int_0^{25} \frac{1}{2}x\,dx = \left[\frac{1}{4}x^2\right]_0^{25} = \frac{625}{4}$ ft-lb. The bottom half of the rope is lifted 25 ft and the work needed to accomplish that is $W_2 = \int_{25}^{50} \frac{1}{2} \cdot 25\,dx = \frac{25}{2}\left[x\right]_{25}^{50} = \frac{625}{2}$ ft-lb. The total work done in pulling half the rope to the top of the building is $W = W_1 + W_2 = \frac{625}{2} + \frac{625}{4} = \frac{3}{4} \cdot 625 = \frac{1875}{4}$ ft-lb.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

14. (a) The 60 ft cable weighs 180 lb, or 3 lb/ft. If we divide the cable into n equal parts of length $\Delta x = 60/n$ ft, then for large n, all points in the ith part are lifted by approximately the same amount. Choose a representative distance from the winch in the ith part of the cable, say x_i^*. If $x_i^* < 25$ ft, then the ith part has to be lifted roughly x_i^* ft. If $x_i^* \geq 25$ ft, then the ith part has to be lifted 25 ft. The ith part weighs $(3 \text{ lb/ft})(\Delta x \text{ ft}) = 3\,\Delta x$ lb, so the work done in lifting it is $(3\,\Delta x)x_i^*$ if $x_i^* < 25$ ft and $(3\,\Delta x)(25) = 75\,\Delta x$ if $x_i^* \geq 25$ ft. The work of lifting the top 25 ft of the cable is $W_1 = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n_1} 3x_i^*\,\Delta x = \int_0^{25} 3x\,dx = \left[\frac{3}{2}x^2\right]_0^{25} = \frac{3}{2}(625) = 937.5$ ft-lb. Here n_1 represents the number of parts of the cable in the top 25 ft. The work of lifting the bottom 35 ft of the cable is $W_2 = \lim\limits_{n\to\infty} \sum\limits_{i=1}^{n_2} 75\,\Delta x = \int_{25}^{60} 75\,dx = 75(60-25) = 2625$ ft-lb, where n_2 represents the number of small parts in the bottom 35 feet of the cable. The total work done is $W = W_1 + W_2 = 937.5 + 2625 = 3562.5$ ft-lb.

(b) Once x feet of cable have been wound up by the winch, there is $(60 - x)$ ft of cable still hanging from the winch. That portion of the cable weighs $3(60 - x)$ lb. Lifting it Δx feet requires $3(60 - x)\,\Delta x$ ft-lb of work. Thus, the total work needed to lift the cable 25 ft is $W = \int_0^{25} 3(60 - x)\,dx = \left[180x - \frac{3}{2}x^2\right]_0^{25} = 4500 - 937.5 = 3562.5$ ft-lb.

15. The work needed to lift the cable is $\lim\limits_{n\to\infty} \sum_{i=1}^{n} 2x_i^*\,\Delta x = \int_0^{500} 2x\,dx = \left[x^2\right]_0^{500} = 250{,}000$ ft-lb. The work needed to lift the coal is 800 lb · 500 ft = 400,000 ft-lb. Thus, the total work required is 250,000 + 400,000 = 650,000 ft-lb.

16. *Assumptions*:

1. After lifting, the chain is L-shaped, with 4 m of the chain lying along the ground.
2. The chain slides effortlessly and without friction along the ground while its end is lifted.
3. The weight density of the chain is constant throughout its length and therefore equals $(8 \text{ kg/m})(9.8 \text{ m/s}^2) = 78.4 \text{ N/m}$.

The part of the chain x m from the lifted end is raised $6 - x$ m if $0 \leq x \leq 6$ m, and it is lifted 0 m if $x > 6$ m. Thus, the work needed is

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} (6 - x_i^*) \cdot 78.4\,\Delta x = \int_0^6 (6 - x)78.4\,dx = 78.4\left[6x - \tfrac{1}{2}x^2\right]_0^6 = (78.4)(18) = 1411.2 \text{ J}$$

17. The chain's weight density is $\dfrac{25 \text{ lb}}{10 \text{ ft}} = 2.5$ lb/ft. The part of the chain x ft below the ceiling (for $5 \leq x \leq 10$) has to be lifted $2(x - 5)$ ft, so the work needed to lift the ith subinterval of the chain is $2(x_i^* - 5)(2.5\,\Delta x)$. The total work needed is

$$\begin{aligned} W &= \lim_{n\to\infty} \sum_{i=1}^{n} 2(x_i^* - 5)(2.5)\,\Delta x = \int_5^{10} [2(x - 5)(2.5)]\,dx = 5\int_5^{10} (x - 5)\,dx \\ &= 5\left[\tfrac{1}{2}x^2 - 5x\right]_5^{10} = 5\left[(50 - 50) - \left(\tfrac{25}{2} - 25\right)\right] = 5\left(\tfrac{25}{2}\right) = 62.5 \text{ ft-lb} \end{aligned}$$

18. The work needed to lift the model rocket itself is $Fd = (mg)d = (0.4 \text{ kg})(9.8 \text{ m/s}^2)(20 \text{ m}) = 78.4$ N-m $= 78.4$ J. At time t (in seconds) the rocket is $x_i^* = 4t$ m above the ground, but it now holds only $(0.75 - 0.15t)$ kg of rocket fuel. In terms

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

of distance, the rocket holds $\left[0.75 - 0.15\left(\frac{1}{4}x_i^*\right)\right]$ kg of rocket fuel when it is x_i^* m above the ground. Moving this mass of rocket fuel a distance of Δx m requires $(9.8)(0.75 - 0.0375x_i^*)\,\Delta x$ J of work. Thus, the work needed to lift the rocket fuel is

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} 9.8(0.75 - 0.0375x_i^*)\,\Delta x = \int_0^{20} 9.8(0.75 - 0.0375x)\,dx$$
$$= 9.8\Big[0.75x - 0.01875x^2\Big]_0^{20} = 9.8[(15 - 7.5) - 0] = 73.5 \text{ J}$$

Adding the work of lifting the model rocket itself gives a total of 151.9 J of work.

19. At a height of x meters ($0 \le x \le 12$), the mass of the rope is $(0.8 \text{ kg/m})(12 - x \text{ m}) = (9.6 - 0.8x)$ kg and the mass of the water is $\left(\frac{36}{12} \text{ kg/m}\right)(12 - x \text{ m}) = (36 - 3x)$ kg. The mass of the bucket is 10 kg, so the total mass is $(9.6 - 0.8x) + (36 - 3x) + 10 = (55.6 - 3.8x)$ kg, and hence, the total force is $9.8(55.6 - 3.8x)$ N. The work needed to lift the bucket Δx m through the ith subinterval of $[0, 12]$ is $9.8(55.6 - 3.8x_i^*)\Delta x$, so the total work is

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} 9.8(55.6 - 3.8x_i^*)\,\Delta x = \int_0^{12} (9.8)(55.6 - 3.8x)\,dx = 9.8\Big[55.6x - 1.9x^2\Big]_0^{12} = 9.8(393.6) \approx 3857 \text{ J}$$

20. A horizontal cylindrical slice of water Δx ft thick has a volume of $\pi r^2 h = \pi \cdot 12^2 \cdot \Delta x$ ft^3 and weighs about $\left(62.5 \text{ lb/ft}^3\right)\left(144\pi\,\Delta x \text{ ft}^3\right) = 9000\pi\,\Delta x$ lb. If the slice lies x_i^* ft below the edge of the pool (where $1 \le x_i^* \le 5$), then the work needed to pump it out is about $9000\pi x_i^*\,\Delta x$. Thus,

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} 9000\pi x_i^*\,\Delta x = \int_1^5 9000\pi x\,dx = \left[4500\pi x^2\right]_1^5 = 4500\pi(25 - 1) = 108{,}000\pi \text{ ft-lb}$$

21. A "slice" of water Δx m thick and lying at a depth of x_i^* m (where $0 \le x_i^* \le \frac{1}{2}$) has volume $(2 \times 1 \times \Delta x)$ m^3, a mass of $2000\,\Delta x$ kg, weighs about $(9.8)(2000\,\Delta x) = 19{,}600\,\Delta x$ N, and thus requires about $19{,}600x_i^*\,\Delta x$ J of work for its removal. So $W = \lim_{n\to\infty} \sum_{i=1}^{n} 19{,}600x_i^*\,\Delta x = \int_0^{1/2} 19{,}600x\,dx = \left[9800x^2\right]_0^{1/2} = 2450$ J.

22. We use a vertical coodinate x measured from the center of the water tank. The top and bottom of the tank have coordinates $x = -12$ ft and $x = 12$ ft, respectively.



A thin horizontal slice of water at coordinate x is a disk of radius $\sqrt{12^2 - x^2}$ as shown in the figure. The disk has area $\pi r^2 = \pi(12^2 - x^2)$, so if the slice has thickness Δx, the slice has volume $\pi(12^2 - x^2)\,\Delta x$ and weight $62.5\pi(12^2 - x^2)\,\Delta x$. The work needed to raise this water from ground level (coordinate 72) to coordinate x, a distance of $(72 - x)$ ft, is $62.5\pi(12^2 - x^2)(72 - x)\,\Delta x$ ft-lb. The total work needed to fill the tank is



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

approximated by a Riemann sum $\sum_{i=1}^{n} 62.5\pi[(12^2-(x_i^*)^2)](72-x_i^*)\,\Delta x$. Thus, the total work is

$$\begin{aligned} W &= \lim_{n\to\infty}\sum_{i=1}^{n} 62.5\pi[(12^2-(x_i^*)^2)](72-x_i^*)\,\Delta x = \int_{-12}^{12} 62.5\pi(12^2-x^2)(72-x)\,dx \\ &= 62.5\pi\int_{-12}^{12}[\underbrace{72(12^2-x^2)}_{\text{even function}} - \underbrace{x(12^2-x^2)}_{\text{odd function}}]\,dx = 62.5\pi(2)\int_0^{12} 72(12^2-x^2)\,dx \qquad \text{[by Theorem 5.5.7]} \\ &= 125\pi(72)\left[12^2x-\tfrac{1}{3}x^3\right]_0^{12} = 9000\pi\left(12^3-\tfrac{1}{3}\cdot 12^3\right) = 9000\pi\left(\tfrac{2}{3}\cdot 12^3\right) \\ &= 10{,}368{,}000\pi \text{ ft-lb} \end{aligned}$$

The 1.5 horsepower pump does $1.5(550) = 825$ ft-lb of work per second. To fill the tank, it will take

$$\frac{10{,}368{,}000\pi \text{ ft-lb}}{825 \text{ ft-lb/s}} \approx 39{,}481 \text{ s} \approx 10.97 \text{ hours}.$$

23. A rectangular "slice" of water Δx m thick and lying x m above the bottom has width x m and volume $8x\,\Delta x$ m^3. It weighs about $(9.8\times 1000)(8x\,\Delta x)$ N, and must be lifted $(5-x)$ m by the pump, so the work needed is about $(9.8\times 10^3)(5-x)(8x\,\Delta x)$ J. The total work required is

$$\begin{aligned} W &\approx \int_0^3 (9.8\times 10^3)(5-x)8x\,dx = (9.8\times 10^3)\int_0^3 (40x-8x^2)\,dx = (9.8\times 10^3)\left[20x^2-\tfrac{8}{3}x^3\right]_0^3 \\ &= (9.8\times 10^3)(180-72) = (9.8\times 10^3)(108) = 1058.4\times 10^3 \approx 1.06\times 10^6 \text{ J} \end{aligned}$$

24. Let y measure depth (in meters) below the center of the spherical tank, so that $y=-3$ at the top of the tank and $y=-4$ at the spigot. A horizontal disk-shaped "slice" of water Δy m thick and lying at coordinate y has radius $\sqrt{9-y^2}$ m and volume $\pi r^2\Delta y = \pi(9-y^2)\,\Delta y$ m^3. It weighs about $(9.8\times 1000)\pi(9-y^2)\,\Delta y$ N and must be lifted $(y+4)$ m by the pump, so the work needed to pump it out is about $(9.8\times 10^3)(y+4)\pi(9-y^2)\,\Delta y$ J. The total work required is

$$\begin{aligned} W &\approx \int_{-3}^{3}(9.8\times 10^3)(y+4)\pi(9-y^2)\,dy = (9.8\times 10^3)\pi\int_{-3}^{3}[y(9-y^2)+4(9-y^2)]\,dy \\ &= (9.8\times 10^3)\pi(2)(4)\int_0^3 (9-y^2)\,dy \qquad \text{[by Theorem 5.5.7]} \\ &= (78.4\times 10^3)\pi\left[9y-\tfrac{1}{3}y^3\right]_0^3 = (78.4\times 10^3)\pi(18) = 1{,}411{,}200\pi \approx 4.43\times 10^6 \text{ J} \end{aligned}$$

25. Let x measure depth (in feet) below the spout at the top of the tank. A horizontal disk-shaped "slice" of water Δx ft thick and lying at coordinate x has radius $\frac{3}{8}(16-x)$ ft $(\star)$ and volume $\pi r^2\Delta x = \pi\cdot\frac{9}{64}(16-x)^2\,\Delta x$ ft^3. It weighs about $(62.5)\frac{9\pi}{64}(16-x)^2\,\Delta x$ lb and must be lifted x ft by the pump, so the work needed to pump it out is about $(62.5)x\,\frac{9\pi}{64}(16-x)^2\,\Delta x$ ft-lb. The total work required is

$$\begin{aligned} W &\approx \int_0^8 (62.5)x\,\tfrac{9\pi}{64}(16-x)^2\,dx = (62.5)\tfrac{9\pi}{64}\int_0^8 x(256-32x+x^2)\,dx \\ &= (62.5)\tfrac{9\pi}{64}\int_0^8 (256x-32x^2+x^3)\,dx = (62.5)\tfrac{9\pi}{64}\left[128x^2-\tfrac{32}{3}x^3+\tfrac{1}{4}x^4\right]_0^8 \\ &= (62.5)\frac{9\pi}{64}\left(\frac{11{,}264}{3}\right) = 33{,}000\pi \approx 1.04\times 10^5 \text{ ft-lb} \end{aligned}$$



$(\star)$ From similar triangles, $\dfrac{d}{8-x} = \dfrac{3}{8}$.

So $r = 3+d = 3+\frac{3}{8}(8-x)$

$= \dfrac{3(8)}{8}+\dfrac{3}{8}(8-x)$

$= \frac{3}{8}(16-x)$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

26. Let x measure the distance (in feet) above the bottom of the tank. A horizontal "slice" of water Δx ft thick and lying at coordinate x has volume $10(2x)\,\Delta x$ ft^3. It weighs about $(62.5)20x\,\Delta x$ lb and must be lifted $(6-x)$ ft by the pump, so the work needed to pump it out is about $(62.5)(6-x)20x\,\Delta x$ ft-lb. The total work required is



$W \approx \int_0^6 (62.5)(6-x)20x\,dx = 1250\int_0^6 (6x-x^2)\,dx = 1250\left[3x^2-\frac{1}{3}x^3\right]_0^6 = 1250(36) = 45{,}000$ ft-lb.

27. If only 4.7×10^5 J of work is done, then only the water above a certain level (call it h) will be pumped out. So we use the same formula as in Exercise 23, except that the work is fixed, and we are trying to find the lower limit of integration:



$4.7\times10^5 \approx \int_h^3 (9.8\times10^3)(5-x)8x\,dx = \left(9.8\times10^3\right)\left[20x^2-\frac{8}{3}x^3\right]_h^3 \quad \Leftrightarrow$

$\frac{4.7}{9.8}\times10^2 \approx 48 = \left(20\cdot3^2-\frac{8}{3}\cdot3^3\right)-\left(20h^2-\frac{8}{3}h^3\right) \quad \Leftrightarrow$

$2h^3-15h^2+45=0$. To find the solution of this equation, we plot $2h^3-15h^2+45$ between $h=0$ and $h=3$. We see that the equation is satisfied for $h\approx2.0$. So the depth of water remaining in the tank is about 2.0 m.

28. The only changes needed in the solution for Exercise 24 are: (1) change the lower limit from -3 to 0 and (2) change 1000 to 900.

$$\begin{aligned} W &\approx \int_0^3 (9.8\times900)(y+4)\pi(9-y^2)\,dy = (9.8\times900)\,\pi\int_0^3 (9y-y^3+36-4y^2)\,dy \\ &= (9.8\times900)\pi\left[\tfrac{9}{2}y^2-\tfrac{1}{4}y^4+36y-\tfrac{4}{3}y^3\right]_0^3 = (9.8\times900)\pi(92.25) = 813{,}645\pi \\ &\approx 2.56\times10^6 \text{ J} \quad \text{[about 58\% of the work in Exercise 24]} \end{aligned}$$

29. $V=\pi r^2x$, so V is a function of x and P can also be regarded as a function of x. If $V_1=\pi r^2x_1$ and $V_2=\pi r^2x_2$, then

$$\begin{aligned} W &= \int_{x_1}^{x_2} F(x)\,dx = \int_{x_1}^{x_2} \pi r^2 P(V(x))\,dx = \int_{x_1}^{x_2} P(V(x))\,dV(x) \qquad [\text{Let } V(x)=\pi r^2x, \text{ so } dV(x)=\pi r^2\,dx.] \\ &= \int_{V_1}^{V_2} P(V)\,dV \quad \text{by the Substitution Rule.} \end{aligned}$$

30. 160 lb/in^2 $=160\cdot144$ lb/ft^2, 100 in^3 $=\frac{100}{1728}$ ft^3, and 800 in^3 $=\frac{800}{1728}$ ft^3.

$k=PV^{1.4}=(160\cdot144)\left(\frac{100}{1728}\right)^{1.4}=23{,}040\left(\frac{25}{432}\right)^{1.4}\approx426.5$. Therefore, $P\approx426.5V^{-1.4}$ and

$$W=\int_{100/1728}^{800/1728} 426.5V^{-1.4}\,dV = 426.5\left[\tfrac{1}{-0.4}V^{-0.4}\right]_{25/432}^{25/54} = (426.5)(2.5)\left[\left(\tfrac{432}{25}\right)^{0.4}-\left(\tfrac{54}{25}\right)^{0.4}\right]\approx1.88\times10^3 \text{ ft-lb.}$$

31.
$$\begin{aligned} W &= \int_{x_1}^{x_2} f(x)\,dx = \int_{t_1}^{t_2} f(s(t))\,v(t)\,dt \quad \left[\begin{matrix} x=s(t), \\ dx=v(t)\,dt \end{matrix}\right] \\ &= \int_{t_1}^{t_2} m\,a(t)\,v(t)\,dt = \int_{v_1}^{v_2} m\,u\,du \quad \left[\begin{matrix} u=v(t), \\ du=a(t)\,dt \end{matrix}\right] \\ &= \left[\tfrac{1}{2}mu^2\right]_{v_1}^{v_2} = \tfrac{1}{2}mv_2^2-\tfrac{1}{2}mv_1^2 \end{aligned}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

32. The mass of the bowling ball is $\dfrac{12 \text{ lb}}{32 \text{ ft/s}^2} = \dfrac{3}{8}$ slug. Converting 20 mi/h to ft/s^2 gives us

$\dfrac{20 \text{ mi}}{\text{h}} \cdot \dfrac{5280 \text{ ft}}{1 \text{ mi}} \cdot \dfrac{1 \text{ h}}{3600 \text{ s}^2} = \dfrac{88}{3}$ ft/s^2. From part (a) with $v_1 = 0$ and $v_2 = \frac{88}{3}$, the work required to hurl the bowling ball is $W = \frac{1}{2} \cdot \frac{3}{8}\left(\frac{88}{3}\right)^2 - \frac{1}{2} \cdot \frac{3}{8}(0)^2 = \frac{484}{3} = 161.\overline{3}$ ft-lb.

33. The work required to move the 800-kg roller coaster car is

$W = \int_0^{60} (5.7x^2 + 1.5x)\,dx = \left[1.9x^3 + 0.75x^2\right]_0^{60} = 410{,}400 + 2700 = 413{,}100$ J.

Using Exercise 31 with $v_1 = 0$, we get $W = \frac{1}{2}mv_2^2 \quad \Rightarrow \quad v_2 = \sqrt{\dfrac{2W}{m}} = \sqrt{\dfrac{2(413{,}100)}{800}} \approx 32.14$ m/s.

34. $W = \int_1^2 \cos\left(\frac{1}{3}\pi x\right) dx = \frac{3}{\pi}\left[\sin\left(\frac{1}{3}\pi x\right)\right]_1^2 = \frac{3}{\pi}\left(\frac{\sqrt{3}}{2} - \frac{\sqrt{3}}{2}\right) = 0$ N·m $= 0$ J.

Interpretation: From $x = 1$ to $x = \frac{3}{2}$, the force does work equal to $\int_1^{3/2} \cos\left(\frac{1}{3}\pi x\right) dx = \frac{3}{\pi}\left(1 - \frac{\sqrt{3}}{2}\right)$ J in accelerating the particle and increasing its kinetic energy. From $x = \frac{3}{2}$ to $x = 2$, the force opposes the motion of the particle, decreasing its kinetic energy. This is negative work, equal in magnitude but opposite in sign to the work done from $x = 1$ to $x = \frac{3}{2}$.

35. (a) $W = \displaystyle\int_a^b F(r)\,dr = \int_a^b G\frac{m_1 m_2}{r^2}\,dr = Gm_1m_2\left[\frac{-1}{r}\right]_a^b = Gm_1m_2\left(\frac{1}{a} - \frac{1}{b}\right)$

(b) By part (a), $W = GMm\left(\dfrac{1}{R} - \dfrac{1}{R + 1{,}000{,}000}\right)$ where $M =$ mass of the earth in kg, $R =$ radius of the earth in m, and $m =$ mass of satellite in kg. (Note that 1000 km $= 1{,}000{,}000$ m.) Thus,

$$W = (6.67 \times 10^{-11})(5.98 \times 10^{24})(1000) \times \left(\frac{1}{6.37 \times 10^6} - \frac{1}{7.37 \times 10^6}\right) \approx 8.50 \times 10^9 \text{ J}$$

36. (a) Assume the pyramid has smooth sides. From the figure for $0 \le x \le 378$, an equation for the side is $y = \frac{-481}{378}x + 481 \quad \Leftrightarrow \quad x = -\frac{378}{481}(y - 481)$. The horizontal length of a cross-section is $2x$ and the area of a cross-section is $A = (2x)^2 = 4x^2 = 4\,\dfrac{378^2}{481^2}(y - 481)^2$. A slice of thickness Δy at height y has volume $\Delta V = A\,\Delta y$ ft^3 and weight $150\,\Delta V$ lb, so the work needed to build the pyramid was



$$W_1 = \int_0^{481} 150y \cdot 4\,\frac{378^2}{481^2}(y - 481)^2\,dy = 600\,\frac{378^2}{481^2}\int_0^{481} (y^3 - 2 \cdot 481y^2 + 481^2 y)\,dy$$

$$= 600\,\frac{378^2}{481^2}\left[\frac{1}{4}y^4 - \frac{2 \cdot 481}{3}y^3 + \frac{481^2}{2}y^2\right]_0^{481} = 600\,\frac{378^2}{481^2}\left(\frac{481^4}{4} - \frac{2 \cdot 481^4}{3} + \frac{481^4}{2}\right)$$

$$= 600\,\frac{378^2}{481^2}\,\frac{481^4}{12} = 50 \cdot 378^2 \cdot 481^2 \approx 1.653 \times 10^{12} \text{ ft-lb}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Work done $= W_2 = \dfrac{10\text{ h}}{\text{day}} \cdot \dfrac{340\text{ days}}{\text{year}} \cdot \dfrac{20\text{ yr}}{1\text{ laborer}} \cdot \dfrac{200\text{ ft-lb}}{\text{hour}} = 1.36 \times 10^7 \dfrac{\text{ft-lb}}{\text{laborer}}$. Dividing W_1 by W_2 gives us about 121,536 laborers.

6.5 Average Value of a Function

1. $f_{\text{avg}} = \frac{1}{b-a}\int_a^b f(x)\,dx = \frac{1}{2-(-1)}\int_{-1}^{2}(3x^2+8x)\,dx = \frac{1}{3}[x^3+4x^2]_{-1}^{2} = \frac{1}{3}[(8+16)-(-1+4)] = 7$

2. $f_{\text{avg}} = \frac{1}{b-a}\int_a^b f(x)\,dx = \frac{1}{4-0}\int_0^4 \sqrt{x}\,dx = \frac{1}{4}\left[\frac{2}{3}x^{3/2}\right]_0^4 = \frac{1}{4}\left(\frac{2}{3}\cdot 8\right) = \frac{4}{3}$

3. $g_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b g(x)\,dx = \frac{1}{\pi/2-(-\pi/2)}\int_{-\pi/2}^{\pi/2} 3\cos x\,dx = \frac{3\cdot 2}{\pi}\int_0^{\pi/2}\cos x\,dx$ [by Theorem 5.5.7(a)]

$= \frac{6}{\pi}\left[\sin x\right]_0^{\pi/2} = \frac{6}{\pi}(1-0) = \frac{6}{\pi}$

4. $f_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b f(z)\,dz = \frac{1}{4-1}\int_1^4 \frac{e^{1/z}}{z^2}\,dz = \frac{1}{3}\int_1^{1/4} e^u\,(-du)$ $\quad\left[\begin{matrix} u = 1/z, \\ du = -1/z^2\,dz \end{matrix}\right]$

$= -\dfrac{1}{3}\left[e^u\right]_1^{1/4} = -\dfrac{1}{3}(e^{1/4}-e) = \dfrac{1}{3}(e-e^{1/4})$

5. $g_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b g(t)\,dt = \frac{1}{2-0}\int_0^2 \frac{9}{1+t^2}\,dt = \frac{1}{2}\left[9\arctan t\right]_0^2 = \frac{9}{2}(\arctan 2 - \arctan 0) = \frac{9}{2}\arctan 2$

6. $f_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b f(x)\,dx = \frac{1}{1-(-1)}\int_{-1}^{1}\frac{x^2}{(x^3+3)^2}\,dx = \frac{1}{2}\int_2^4 \frac{1}{u^2}\left(\frac{1}{3}\,du\right)$ $\quad\left[\begin{matrix} u = x^3+3, \\ du = 3x^2\,dx \end{matrix}\right]$

$= \dfrac{1}{6}\left[-\dfrac{1}{u}\right]_2^4 = \dfrac{1}{6}\left(-\dfrac{1}{4}+\dfrac{1}{2}\right) = \dfrac{1}{24}$

7. $h_{\text{avg}} = \frac{1}{b-a}\int_a^b h(x)\,dx = \frac{1}{\pi-0}\int_0^{\pi}\cos^4 x\,\sin x\,dx = \frac{1}{\pi}\int_1^{-1} u^4(-du)$ $\quad[u=\cos x,\, du = -\sin x\,dx]$

$= \frac{1}{\pi}\int_{-1}^{1} u^4\,du = \frac{1}{\pi}\cdot 2\int_0^1 u^4\,du$ [by Theorem 5.5.7(a)] $= \frac{2}{\pi}\left[\frac{1}{5}u^5\right]_0^1 = \frac{2}{5\pi}$

8. $h_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b h(u)\,du = \frac{1}{5-1}\int_1^5 \frac{\ln u}{u}\,du = \frac{1}{4}\int_0^{\ln 5} y\,dy$ $\quad\left[\begin{matrix} y = \ln u, \\ dy = 1/u\,du \end{matrix}\right]$

$= \frac{1}{4}\left[\frac{1}{2}y^2\right]_0^{\ln 5} = \frac{1}{8}(\ln 5)^2$

9. (a) $f_{\text{avg}} = \dfrac{1}{3-1}\displaystyle\int_1^3 \frac{1}{t^2}\,dt = \frac{1}{2}\left[-\frac{1}{t}\right]_1^3$

$= \dfrac{1}{2}\left[-\dfrac{1}{3}-(-1)\right] = \dfrac{1}{3}$

(b) $f(c) = f_{\text{avg}} \;\Leftrightarrow\; \dfrac{1}{c^2} = \dfrac{1}{3} \;\Leftrightarrow\; c^2 = 3 \;\Rightarrow$

$c = \sqrt{3}$ since $-\sqrt{3}$ is not in $[1,3]$.

(c)



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

10. (a) $g_{\text{avg}} = \dfrac{1}{2-0}\displaystyle\int_0^2 (x+1)^3\,dx = \frac{1}{2}\left[\frac{1}{4}(x+1)^4\right]_0^2$

$= \dfrac{1}{8}(3^4 - 1^4) = \dfrac{1}{8}\cdot 80 = 10$

(b) $g(c) = g_{\text{avg}} \;\Leftrightarrow\; (c+1)^3 = 10 \;\Leftrightarrow\; c+1 = \sqrt[3]{10}$

$\Leftrightarrow\; c = -1 + \sqrt[3]{10} \approx 1.154$

(c)



11. (a) $f_{\text{avg}} = \dfrac{1}{\pi - 0}\displaystyle\int_0^{\pi} (2\sin x - \sin 2x)\,dx$

$= \frac{1}{\pi}\left[-2\cos x + \frac{1}{2}\cos 2x\right]_0^{\pi}$

$= \frac{1}{\pi}\left[\left(2 + \frac{1}{2}\right) - \left(-2 + \frac{1}{2}\right)\right] = \frac{4}{\pi}$

(b) $f(c) = f_{\text{avg}} \;\Leftrightarrow\; 2\sin c - \sin 2c = \frac{4}{\pi} \;\Leftrightarrow$

$c = c_1 \approx 1.238$ or $c = c_2 \approx 2.808$

(c)



12. (a) $f_{\text{avg}} = \frac{1}{2-0}\int_0^2 2xe^{-x^2}\,dx$

$= \frac{1}{2}\left[-e^{-x^2}\right]_0^2 = \frac{1}{2}(-e^{-4} + 1)$

(b) $f(c) = f_{\text{avg}} \;\Leftrightarrow\; 2ce^{-c^2} = \frac{1}{2}(1 - e^{-4}) \;\Leftrightarrow$

$c = c_1 \approx 0.263$ or $c = c_2 \approx 1.287$

(c)



13. f is continuous on $[1,3]$, so by the Mean Value Theorem for Integrals there exists a number c in $[1,3]$ such that $\int_1^3 f(x)\,dx = f(c)(3-1) \;\Rightarrow\; 8 = 2f(c)$; that is, there is a number c such that $f(c) = \frac{8}{2} = 4$.

14. The requirement is that $\dfrac{1}{b-0}\displaystyle\int_0^b f(x)\,dx = 3$. The LHS of this equation is equal to

$\dfrac{1}{b}\displaystyle\int_0^b \left(2 + 6x - 3x^2\right)dx = \frac{1}{b}\left[2x + 3x^2 - x^3\right]_0^b = 2 + 3b - b^2$, so we solve the equation $2 + 3b - b^2 = 3 \;\Leftrightarrow$

$b^2 - 3b + 1 = 0 \;\Leftrightarrow\; b = \dfrac{3 \pm \sqrt{(-3)^2 - 4\cdot 1\cdot 1}}{2\cdot 1} = \dfrac{3 \pm \sqrt{5}}{2}$. Both roots are valid since they are positive.

15. Use geometric interpretations to find the values of the integrals.

$\int_0^8 f(x)\,dx = \int_0^1 f(x)\,dx + \int_1^2 f(x)\,dx + \int_2^3 f(x)\,dx + \int_3^4 f(x)\,dx + \int_4^6 f(x)\,dx + \int_6^7 f(x)\,dx + \int_7^8 f(x)\,dx$

$= -\frac{1}{2} + \frac{1}{2} + \frac{1}{2} + 1 + 4 + \frac{3}{2} + 2 = 9$

Thus, the average value of f on $[0,8] = f_{\text{avg}} = \frac{1}{8-0}\int_0^8 f(x)\,dx = \frac{1}{8}(9) = \frac{9}{8}$.

16. (a) $v_{\text{avg}} = \frac{1}{12-0}\int_0^{12} v(t)\,dt = \frac{1}{12}I$. Use the Midpoint Rule with $n = 3$ and $\Delta t = \frac{12-0}{3} = 4$ to estimate I.

$I \approx M_3 = 4[v(2) + v(6) + v(10)] = 4[21 + 50 + 66] = 4(137) = 548$. Thus, $v_{\text{avg}} \approx \frac{1}{12}(548) = 45\frac{2}{3}$ km/h.

(b) Estimating from the graph, $v(t) = 45\frac{2}{3}$ when $t \approx 5.2$ s.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

17. Let $t = 0$ and $t = 12$ correspond to 9 AM and 9 PM, respectively.

$$T_{\text{avg}} = \tfrac{1}{12-0}\int_0^{12}\left[50 + 14\sin\tfrac{1}{12}\pi t\right]dt = \tfrac{1}{12}\left[50t - 14\cdot\tfrac{12}{\pi}\cos\tfrac{1}{12}\pi t\right]_0^{12}$$
$$= \tfrac{1}{12}\left[50\cdot 12 + 14\cdot\tfrac{12}{\pi} + 14\cdot\tfrac{12}{\pi}\right] = 50 + \tfrac{28}{\pi} \approx 59\,^\circ\text{F}$$

18. East Coast: $T_{E_\text{avg}} = \frac{1}{24-0}\int_0^{24} T_E(t)\,dt = \frac{1}{24}E$; West Coast: $T_{W_\text{avg}} = \frac{1}{24-0}\int_0^{24} T_W(t)\,dt = \frac{1}{24}W$

Use the Midpoint Rule with $n = 12$ and $\Delta t = \frac{24-0}{12} = 2$ to estimate E and W.

$$E \approx M_{12} = 2[T_E(1) + T_E(3) + \cdots + T_E(23)] = 2(60 + 60 + 65 + 65 + 75 + 70 + 65 + 70 + 55 + 65 + 60 + 55)$$
$$= 2(765) = 1530$$

$$W \approx M_{12} = 2[T_W(1) + T_W(3) + \cdots + T_W(23)] = 2(65 + 55 + 50 + 55 + 60 + 65 + 80 + 65 + 60 + 55 + 50 + 50)$$
$$= 2(710) = 1420$$

Thus, $T_{E_\text{avg}} = \frac{1}{24}(1530) = 63\frac{3}{4}\,^\circ\text{F}$, and $T_{W_\text{avg}} = \frac{1}{24}(1420) = 59\frac{1}{6}\,^\circ\text{F}$.

While the West Coast city had the highest temperature that day, 80 °F, the East Coast city was "warmer" overall, because the East Coast city had the higher average temperature.

19. $\rho_{\text{avg}} = \dfrac{1}{8}\displaystyle\int_0^8 \frac{12}{\sqrt{x+1}}\,dx = \frac{3}{2}\int_0^8 (x+1)^{-1/2}\,dx = \left[3\sqrt{x+1}\right]_0^8 = 9 - 3 = 6\ \text{kg/m}$

20. $v_{\text{avg}} = \dfrac{1}{R-0}\displaystyle\int_0^R v(r)\,dr = \frac{1}{R}\int_0^R \frac{P}{4\eta l}(R^2 - r^2)\,dr = \frac{P}{4\eta l R}\left[R^2 r - \tfrac{1}{3}r^3\right]_0^R = \frac{P}{4\eta l R}\left(\tfrac{2}{3}\right)R^3 = \frac{PR^2}{6\eta l}.$

Since $v(r)$ is decreasing on $(0, R]$, $v_{\max} = v(0) = \dfrac{PR^2}{4\eta l}$. Thus, $v_{\text{avg}} = \frac{2}{3}v_{\max}$.

21. $P_{\text{avg}} = \frac{1}{50-0}\int_0^{50} P(t)\,dt = \frac{1}{50}\int_0^{50} 2560e^{bt}\,dt \qquad [\text{with } b = 0.017185]$

$$= \frac{2560}{50}\left[\frac{1}{b}e^{bt}\right]_0^{50} = \frac{2560}{50b}(e^{50b} - 1) \approx 4056 \text{ million, or about 4 billion people}$$

22. (a) Similar to Example 3.8.3, we have $T_s = 20°\text{C}$ and hence $\dfrac{dT}{dt} = c(T - 20)$. Let $y = T - 20$, so that

$y(0) = T(0) - 20 = 95 - 20 = 75$. Now y satisfies (3.8.2), so $y = 75e^{ct}$. We are given that $T(30) = 61$, so

$y(30) = 61 - 20 = 41$ and $41 = 75e^{c(30)} \quad\Rightarrow\quad \frac{41}{75} = e^{30c} \quad\Rightarrow\quad 30c = \ln\frac{41}{75} \quad\Rightarrow\quad c = \frac{1}{30}\ln\frac{41}{75} \approx -0.020131.$

Thus, $T(t) = 20 + 75e^{-kt}$, where $k = -c \approx 0.02$.

(b) $T_{\text{avg}} = \frac{1}{30-0}\int_0^{30} T(t)\,dt = \frac{1}{30}\int_0^{30}(20 + 75e^{-kt})\,dt = \frac{1}{30}\left[20t - \frac{75}{k}e^{-kt}\right]_0^{30} = \frac{1}{30}\left[\left(600 - \frac{75}{k}e^{-30k}\right) - \left(0 - \frac{75}{k}\right)\right]$

$$= \tfrac{1}{30}\left(600 - \tfrac{75}{k}\cdot\tfrac{41}{75} + \tfrac{75}{k}\right) = \tfrac{1}{30}\left(600 + \tfrac{34}{k}\right) = 20 + \tfrac{34}{30k} \approx 76.3°\text{C}$$

23. $V_{\text{avg}} = \frac{1}{5}\int_0^5 V(t)\,dt = \frac{1}{5}\int_0^5 \frac{5}{4\pi}\left[1 - \cos\left(\tfrac{2}{5}\pi t\right)\right]dt = \frac{1}{4\pi}\int_0^5 \left[1 - \cos\left(\tfrac{2}{5}\pi t\right)\right]dt$

$$= \tfrac{1}{4\pi}\left[t - \tfrac{5}{2\pi}\sin\left(\tfrac{2}{5}\pi t\right)\right]_0^5 = \tfrac{1}{4\pi}\left[(5 - 0) - 0\right] = \tfrac{5}{4\pi} \approx 0.4\ \text{L}$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

24. $s=\frac{1}{2}gt^2 \;\Rightarrow\; t=\sqrt{2s/g}$ [since $t\ge 0$]. Now $v=ds/dt=gt=g\sqrt{2s/g}=\sqrt{2gs} \;\Rightarrow\; v^2=2gs \;\Rightarrow\; s=\dfrac{v^2}{2g}$.

We see that v can be regarded as a function of t or of s: $v=F(t)=gt$ and $v=G(s)=\sqrt{2gs}$. Note that $v_T=F(T)=gT$.

Displacement can be viewed as a function of t: $s=s(t)=\frac{1}{2}gt^2$; also $s(t)=\dfrac{v^2}{2g}=\dfrac{[F(t)]^2}{2g}$. When $t=T$, these two formulas for $s(t)$ imply that

$$\sqrt{2gs(T)}=F(T)=v_T=gT=2\left(\tfrac{1}{2}gT^2\right)/T=2s(T)/T \qquad (\star)$$

The average of the velocities with respect to time t during the interval $[0,T]$ is

$$v_{t\text{-avg}}=f_{\text{avg}}=\frac{1}{T-0}\int_0^T F(t)\,dt=\frac{1}{T}\left[s(T)-s(0)\right] \quad [\text{by FTC}] \quad =\frac{s(T)}{T} \quad [\text{since } s(0)=0] \quad =\frac{1}{2}v_T \quad [\text{by } (\star)]$$

But the average of the velocities with respect to displacement s during the corresponding displacement interval $[s(0),s(T)]=[0,s(T)]$ is

$$\begin{aligned} v_{s\text{-avg}}=g_{\text{avg}}&=\frac{1}{s(T)-0}\int_0^{s(T)} G(s)\,ds=\frac{1}{s(T)}\int_0^{s(T)}\sqrt{2gs}\,ds=\frac{\sqrt{2g}}{s(T)}\int_0^{s(T)} s^{1/2}\,ds \\ &=\frac{\sqrt{2g}}{s(T)}\cdot\frac{2}{3}\left[s^{3/2}\right]_0^{s(T)}=\frac{2}{3}\cdot\frac{\sqrt{2g}}{s(T)}\cdot\left[s(T)\right]^{3/2}=\frac{2}{3}\sqrt{2gs(T)}=\frac{2}{3}v_T \quad [\text{by } (\star)] \end{aligned}$$

25. $f_{\text{avg}}=\frac{1}{b-a}\int_a^b f(x)\,dx$

$>\frac{1}{b-a}$ (area of trapezoid $ABDF$)

$=\frac{1}{b-a}$ (area of rectangle $ACEF$)

$=\frac{1}{b-a}\left[f\left(\frac{a+b}{2}\right)\cdot(b-a)\right]$

$=f\left(\frac{a+b}{2}\right)$



26. $f_{\text{avg}}\,[a,b]=\dfrac{1}{b-a}\displaystyle\int_a^b f(x)\,dx=\frac{1}{b-a}\int_a^c f(x)\,dx+\frac{1}{b-a}\int_c^b f(x)\,dx$

$$=\frac{c-a}{b-a}\left[\frac{1}{c-a}\int_a^c f(x)\,dx\right]+\frac{b-c}{b-a}\left[\frac{1}{b-c}\int_c^b f(x)\,dx\right]=\frac{c-a}{b-a}\,f_{\text{avg}}\,[a,c]+\frac{b-c}{b-a}\,f_{\text{avg}}\,[c,b]$$

27. Since f is continuous, we can apply l'Hospital's Rule.

$$\lim_{t\to a^+} f_{\text{avg}}[a,t]=\lim_{t\to a^+}\frac{1}{t-a}\int_a^t f(x)\,dx=\lim_{t\to a^+}\frac{\displaystyle\int_a^t f(x)\,dx}{t-a}\ \left[\text{form }\tfrac{0}{0}\right]\overset{\text{H}}{=}\lim_{t\to a^+}\frac{f(t)}{1}=f(a)$$

28. Let $F(x)=\int_a^x f(t)\,dt$ for x in $[a,b]$. Then F is continuous on $[a,b]$ and differentiable on (a,b), so by the Mean Value Theorem there is a number c in (a,b) such that $F(b)-F(a)=F'(c)(b-a)$. But $F'(x)=f(x)$ by the Fundamental Theorem of Calculus. Therefore, $\int_a^b f(t)\,dt-0=f(c)(b-a)$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

APPLIED PROJECT Calculus and Baseball

1. (a) $F = ma = m\,\dfrac{dv}{dt}$, so by the Substitution Rule we have

$$\int_{t_0}^{t_1} F(t)\,dt = \int_{t_0}^{t_1} m\left(\frac{dv}{dt}\right) dt = m\int_{v_0}^{v_1} dv = \big[mv\big]_{v_0}^{v_1} = mv_1 - mv_0 = p(t_1) - p(t_0)$$

(b) (i) We have $v_1 = 110$ mi/h $= \frac{110(5280)}{3600}$ ft/s $= 161.\overline{3}$ ft/s, $v_0 = -90$ mi/h $= -132$ ft/s, and the mass of the baseball is $m = \dfrac{w}{g} = \frac{5/16}{32} = \frac{5}{512}$. So the change in momentum is

$p(t_1) - p(t_0) = mv_1 - mv_0 = \frac{5}{512}\,[161.\overline{3} - (-132)] \approx 2.86$ slug-ft/s.

(ii) From part (a) and part (b)(i), we have $\int_0^{0.001} F(t)\,dt = p(0.001) - p(0) \approx 2.86$, so the average force over the interval $[0, 0.001]$ is $\frac{1}{0.001}\int_0^{0.001} F(t)\,dt \approx \frac{1}{0.001}(2.86) = 2860$ lb.

2. (a) $W = \displaystyle\int_{s_0}^{s_1} F(s)\,ds$, where $F(s) = m\,\dfrac{dv}{dt} = m\,\dfrac{dv}{ds}\dfrac{ds}{dt} = mv\,\dfrac{dv}{ds}$ and so, by the Substitution Rule,

$$W = \int_{s_0}^{s_1} F(s)\,ds = \int_{s_0}^{s_1} mv\,\frac{dv}{ds}\,ds = \int_{v(s_0)}^{v(s_1)} mv\,dv = \big[\tfrac{1}{2}mv^2\big]_{v_0}^{v_1} = \tfrac{1}{2}mv_1^2 - \tfrac{1}{2}mv_0^2$$

(b) From part (b)(i), 90 mi/h $= 132$ ft/s. Assume $v_0 = v(s_0) = 0$ and $v_1 = v(s_1) = 132$ ft/s [note that s_1 is the point of release of the baseball]. $m = \frac{5}{512}$, so the work done is $W = \frac{1}{2}mv_1^2 - \frac{1}{2}mv_0^2 = \frac{1}{2}\cdot\frac{5}{512}\cdot(132)^2 \approx 85$ ft-lb.

3. (a) Here we have a differential equation of the form $dv/dt = kv$, so by Theorem 3.8.2, the solution is $v(t) = v(0)e^{kt}$. In this case $k = -\frac{1}{10}$ and $v(0) = 100$ ft/s, so $v(t) = 100e^{-t/10}$. We are interested in the time t that the ball takes to travel 280 ft, so we find the distance function

$$s(t) = \textstyle\int_0^t v(x)\,dx = \int_0^t 100e^{-x/10}\,dx = 100\Big[-10e^{-x/10}\Big]_0^t = -1000(e^{-t/10} - 1) = 1000(1 - e^{-t/10})$$

Now we set $s(t) = 280$ and solve for t: $280 = 1000(1 - e^{-t/10}) \;\Rightarrow\; 1 - e^{-t/10} = \frac{7}{25} \;\Rightarrow$

$-\frac{1}{10}t = \ln\big(1 - \frac{7}{25}\big) \;\Rightarrow\; t \approx 3.285$ seconds.

(b) Let x be the distance of the shortstop from home plate. We calculate the time for the ball to reach home plate as a function of x, then differentiate with respect to x to find the value of x which corresponds to the minimum time. The total time that it takes the ball to reach home is the sum of the times of the two throws, plus the relay time $\big(\frac{1}{2}\text{ s}\big)$. The distance from the fielder to the shortstop is $280 - x$, so to find the time t_1 taken by the first throw, we solve the equation

$s_1(t_1) = 280 - x \;\Leftrightarrow\; 1 - e^{-t_1/10} = \dfrac{280 - x}{1000} \;\Leftrightarrow\; t_1 = -10\ln\dfrac{720 + x}{1000}$. We find the time t_2 taken by the second throw if the shortstop throws with velocity w, since we see that this velocity varies in the rest of the problem. We use

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$v = we^{-t/10}$ and isolate t_2 in the equation $s(t_2) = 10w(1 - e^{-t_2/10}) = x \quad \Leftrightarrow \quad e^{-t_2/10} = 1 - \dfrac{x}{10w} \quad \Leftrightarrow$

$t_2 = -10 \ln \dfrac{10w - x}{10w}$, so the total time is $t_w(x) = \dfrac{1}{2} - 10\left[\ln \dfrac{720 + x}{1000} + \ln \dfrac{10w - x}{10w}\right]$.

To find the minimum, we differentiate: $\dfrac{dt_w}{dx} = -10\left[\dfrac{1}{720 + x} - \dfrac{1}{10w - x}\right]$, which changes from negative to positive when $720 + x = 10w - x \quad \Leftrightarrow \quad x = 5w - 360$. By the First Derivative Test, t_w has a minimum at this distance from the shortstop to home plate. So if the shortstop throws at $w = 105$ ft/s from a point $x = 5(105) - 360 = 165$ ft from home plate, the minimum time is $t_{105}(165) = \frac{1}{2} - 10\big(\ln \frac{720+165}{1000} + \ln \frac{1050 - 165}{1050}\big) \approx 3.431$ seconds. This is longer than the time taken in part (a), so in this case the manager should encourage a direct throw. If $w = 115$ ft/s, then $x = 215$ ft from home, and the minimum time is $t_{115}(215) = \frac{1}{2} - 10\big(\ln \frac{720+215}{1000} + \ln \frac{1150 - 215}{1150}\big) \approx 3.242$ seconds. This is less than the time taken in part (a), so in this case, the manager should encourage a relayed throw.

(c) In general, the minimum time is $t_w(5w - 360) = \dfrac{1}{2} - 10\left[\ln \dfrac{360 + 5w}{1000} + \ln \dfrac{360 + 5w}{10w}\right] = \dfrac{1}{2} - 10 \ln \dfrac{(w + 72)^2}{400w}$.

We want to find out when this is about 3.285 seconds, the same time as the direct throw. From the graph, we estimate that this is the case for $w \approx 112.8$ ft/s. So if the shortstop can throw the ball with this velocity, then a relayed throw takes the same time as a direct throw.



APPLIED PROJECT Where to Sit at the Movies

1. $|VP| = 9 + x\cos\alpha$, $|PT| = 35 - (4 + x\sin\alpha) = 31 - x\sin\alpha$, and $|PB| = (4 + x\sin\alpha) - 10 = x\sin\alpha - 6$. So using the Pythagorean Theorem, we have $|VT| = \sqrt{|VP|^2 + |PT|^2} = \sqrt{(9 + x\cos\alpha)^2 + (31 - x\sin\alpha)^2} = a$, and $|VB| = \sqrt{|VP|^2 + |PB|^2} = \sqrt{(9 + x\cos\alpha)^2 + (x\sin\alpha - 6)^2} = b$.



Using the Law of Cosines on $\triangle VBT$, we get $25^2 = a^2 + b^2 - 2ab\cos\theta \quad \Leftrightarrow \quad \cos\theta = \dfrac{a^2 + b^2 - 625}{2ab} \quad \Leftrightarrow$

$\theta = \arccos\left(\dfrac{a^2 + b^2 - 625}{2ab}\right)$, as required.

2. From the graph of θ, it appears that the value of x which maximizes θ is $x \approx 8.25$ ft. Assuming that the first row is at $x = 0$, the row closest to this value of x is the fourth row, at $x = 9$ ft, and from the graph, the viewing angle in this row seems to be about 0.85 radians, or about 49°.



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

3. With a CAS, we type in the definition of θ, substitute in the proper values of a and b in terms of x and $\alpha = 20^\circ = \frac{\pi}{9}$ radians, and then use the differentiation command to find the derivative. We use a numerical rootfinder and find that the root of the equation $d\theta/dx = 0$ is $x \approx 8.253062$, as approximated in Problem 2.

4. From the graph in Problem 2, it seems that the average value of the function on the interval $[0, 60]$ is about 0.6. We can use a CAS to approximate $\frac{1}{60}\int_0^{60} \theta(x)\,dx \approx 0.625 \approx 36^\circ$. (The calculation is much faster if we reduce the number of digits of accuracy required.) The minimum value is $\theta(60) \approx 0.38$ and, from Problem 2, the maximum value is about 0.85.

6 Review

TRUE-FALSE QUIZ

1. False. For example, let $f(x) = x$, $g(x) = 2x$, $a = 1$, and $b = 2$. The area between the curves for $a \le x \le b$ is $A = \int_a^b [g(x) - f(x)]\,dx$. The given integral represents area when $f(x) \ge g(x)$ for $a \le x \le b$.

2. False. In a solid of revolution, cross-sections perpendicular to the axis of rotation are circular. A cube has no circular cross-sections.

3. False. Cross-sections perpendicular to the x-axis are washers, and we find cross-sectional area by subtracting the area of the inner circle from the area of the outer circle. Thus, $A(x) = \pi\left(\sqrt{x}\right)^2 - \pi(x)^2 = \pi\left[\left(\sqrt{x}\right)^2 - x^2\right]$, and the volume of the resulting solid is $V = \displaystyle\int_0^1 A(x)\,dx = \int_0^1 \pi\left[\left(\sqrt{x}\right)^2 - x^2\right]\,dx$.

4. True. See "The Method of Cylindrical Shells" in Section 6.3.

5. True. See "Volumes of Solids of Revolution" in Section 6.2.

6. True. See "Volumes of Solids of Revolution" in Section 6.2.

7. False. Cross-sections perpendicular to the y-axis are washers.

8. False. Using the method of cylindrical shells, a typical shell of the solid obtained by revolving $\mathcal{R}$ about the y-axis has radius x, and the volume of the solid is $V = \int_a^b 2\pi x\, f(x)\,dx$. Again using the method of cylindrical shells, a typical shell of the solid obtained by revolving about the line $x = -2$ has radius $x - (-2) = x + 2$, and the volume of the solid is $V = \int_a^b 2\pi(x + 2)\, f(x)\,dx$.

9. True. A cross-section of S perpendicular to the x-axis is a square with side length $f(x)$, so each cross-section has area $A(x) = [f(x)]^2$ and volume $V = \int_a^b A(x)\,dx = \int_a^b [f(x)]^2\,dx$.

10. False. The work done to pull up the top half of the cable will be more than half of the work required to pull up the entire cable. When the top half of the cable is being pulled up, the bottom half is still attached, and extra work is required to pull that bottom half up as the top half is pulled up. There is no such extra work required as the remaining bottom half is pulled up.

11. True. By defnition of the average value of f on the interval $[a, b]$, the average value of f on $[2, 5]$ is $\frac{1}{5-2}\int_2^5 f(x)\,dx = \frac{1}{3}(12) = 4$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

EXERCISES

1. The curves intersect when $x^2 = 8x - x^2 \;\Leftrightarrow\; 2x^2 - 8x = 0 \;\Leftrightarrow\; 2x(x-4) = 0 \;\Leftrightarrow\; x = 0$ or 4.

$$A = \int_0^4 \left[(8x - x^2) - x^2\right] dx = \int_0^4 (8x - 2x^2)\, dx$$
$$= \left[4x^2 - \tfrac{2}{3}x^3\right]_0^4 = \left[\left(64 - \tfrac{128}{3}\right) - 0\right] = \tfrac{64}{3}$$



2. The line $y = x - 2$ intersects the curve $y = \sqrt{x}$ at $(4, 2)$ and it intersects the curve $y = -\sqrt[3]{x}$ at $(1, -1)$.

$$A = \int_0^1 [\sqrt{x} - (-\sqrt[3]{x})]\, dx + \int_1^4 [\sqrt{x} - (x-2)]\, dx$$
$$= \left[\tfrac{2}{3}x^{3/2} + \tfrac{3}{4}x^{4/3}\right]_0^1 + \left[\tfrac{2}{3}x^{3/2} - \tfrac{1}{2}x^2 + 2x\right]_1^4$$
$$= \left(\tfrac{2}{3} + \tfrac{3}{4}\right) - 0 + \left(\tfrac{16}{3} - 8 + 8\right) - \left(\tfrac{2}{3} - \tfrac{1}{2} + 2\right)$$
$$= \tfrac{16}{3} + \tfrac{3}{4} - \tfrac{3}{2} = \tfrac{55}{12}$$

Or, integrating with respect to y: $A = \int_{-1}^0 [(y+2) - (-y^3)]\, dy + \int_0^2 [(y+2) - y^2]\, dy$



3. If $x \geq 0$, then $|x| = x$, and the graphs intersect when $x = 1 - 2x^2 \;\Leftrightarrow\; 2x^2 + x - 1 = 0 \;\Leftrightarrow\; (2x-1)(x+1) = 0 \;\Leftrightarrow\; x = \frac{1}{2}$ or -1, but $-1 < 0$. By symmetry, we can double the area from $x = 0$ to $x = \frac{1}{2}$.

$$A = 2\int_0^{1/2} \left[(1 - 2x^2) - x\right] dx = 2\int_0^{1/2} (-2x^2 - x + 1)\, dx$$
$$= 2\left[-\tfrac{2}{3}x^3 - \tfrac{1}{2}x^2 + x\right]_0^{1/2} = 2\left[\left(-\tfrac{1}{12} - \tfrac{1}{8} + \tfrac{1}{2}\right) - 0\right]$$
$$= 2\left(\tfrac{7}{24}\right) = \tfrac{7}{12}$$



4. $y^2 + 3y = -y \;\Leftrightarrow\; y^2 + 4y = 0 \;\Leftrightarrow\; y(y+4) = 0 \;\Leftrightarrow\; y = 0$ or -4.

$$A = \int_{-4}^0 \left[-y - (y^2 + 3y)\right] dy = \int_{-4}^0 (-y^2 - 4y)\, dy$$
$$= \left[-\tfrac{1}{3}y^3 - 2y^2\right]_{-4}^0 = 0 - \left(\tfrac{64}{3} - 32\right) = \tfrac{32}{3}$$



5. $$A = \int_0^2 \left[\sin\left(\frac{\pi x}{2}\right) - (x^2 - 2x)\right] dx$$
$$= \left[-\frac{2}{\pi}\cos\left(\frac{\pi x}{2}\right) - \frac{1}{3}x^3 + x^2\right]_0^2$$
$$= \left(\tfrac{2}{\pi} - \tfrac{8}{3} + 4\right) - \left(-\tfrac{2}{\pi} - 0 + 0\right) = \tfrac{4}{3} + \tfrac{4}{\pi}$$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

6. $A = \int_0^1 \left(\sqrt{x} - x^2\right) dx + \int_1^2 \left(x^2 - \sqrt{x}\right) dx$

$= \left[\frac{2}{3}x^{3/2} - \frac{1}{3}x^3\right]_0^1 + \left[\frac{1}{3}x^3 - \frac{2}{3}x^{3/2}\right]_1^2$

$= \left[\left(\frac{2}{3} - \frac{1}{3}\right) - 0\right] + \left[\left(\frac{8}{3} - \frac{4}{3}\sqrt{2}\right) - \left(\frac{1}{3} - \frac{2}{3}\right)\right]$

$= \frac{10}{3} - \frac{4}{3}\sqrt{2}$



7. Using washers with inner radius x^2 and outer radius $2x$, we have

$V = \pi \int_0^2 \left[(2x)^2 - (x^2)^2\right] dx = \pi \int_0^2 (4x^2 - x^4)\, dx$

$= \pi\left[\frac{4}{3}x^3 - \frac{1}{5}x^5\right]_0^2 = \pi\left(\frac{32}{3} - \frac{32}{5}\right)$

$= 32\pi \cdot \frac{2}{15} = \frac{64}{15}\pi$



8. $1 + y^2 = y + 3 \quad \Leftrightarrow \quad y^2 - y - 2 = 0 \quad \Leftrightarrow \quad (y - 2)(y + 1) = 0 \quad \Leftrightarrow$

$y = 2$ or -1.

$V = \pi \int_{-1}^2 \left[(y + 3)^2 - (1 + y^2)^2\right] dy = \pi \int_{-1}^2 (y^2 + 6y + 9 - 1 - 2y^2 - y^4)\, dy$

$= \pi \int_{-1}^2 (8 + 6y - y^2 - y^4)\, dy = \pi\left[8y + 3y^2 - \frac{1}{3}y^3 - \frac{1}{5}y^5\right]_{-1}^2$

$= \pi\left[\left(16 + 12 - \frac{8}{3} - \frac{32}{5}\right) - \left(-8 + 3 + \frac{1}{3} + \frac{1}{5}\right)\right] = \pi\left(33 - \frac{9}{3} - \frac{33}{5}\right) = \frac{117}{5}\pi$



9. $V = \pi \int_{-3}^3 \left\{\left[(9 - y^2) - (-1)\right]^2 - \left[0 - (-1)\right]^2\right\} dy$

$= 2\pi \int_0^3 \left[(10 - y^2)^2 - 1\right] dy = 2\pi \int_0^3 (100 - 20y^2 + y^4 - 1)\, dy$

$= 2\pi \int_0^3 (99 - 20y^2 + y^4)\, dy = 2\pi\left[99y - \frac{20}{3}y^3 + \frac{1}{5}y^5\right]_0^3$

$= 2\pi\left(297 - 180 + \frac{243}{5}\right) = \frac{1656}{5}\pi$



10. $V = \pi \int_{-2}^2 \left\{\left[(9 - x^2) - (-1)\right]^2 - \left[(x^2 + 1) - (-1)\right]^2\right\} dx$

$= \pi \int_{-2}^2 \left[(10 - x^2)^2 - (x^2 + 2)^2\right] dx$

$= 2\pi \int_0^2 (96 - 24x^2)\, dx = 48\pi \int_0^2 (4 - x^2)\, dx$

$= 48\pi\left[4x - \frac{1}{3}x^3\right]_0^2 = 48\pi\left(8 - \frac{8}{3}\right) = 256\pi$



© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

11. The graph of $x^2 - y^2 = a^2$ is a hyperbola with right and left branches.



Solving for y gives us $y^2 = x^2 - a^2 \quad \Rightarrow \quad y = \pm\sqrt{x^2 - a^2}$.

We'll use shells and the height of each shell is

$\sqrt{x^2 - a^2} - \left(-\sqrt{x^2 - a^2}\right) = 2\sqrt{x^2 - a^2}$.

The volume is $V = \int_a^{a+h} 2\pi x \cdot 2\sqrt{x^2 - a^2}\, dx$. To evaluate, let $u = x^2 - a^2$, so $du = 2x\, dx$ and $x\, dx = \frac{1}{2}\, du$. When $x = a$, $u = 0$, and when $x = a + h$, $u = (a + h)^2 - a^2 = a^2 + 2ah + h^2 - a^2 = 2ah + h^2$.

Thus, $V = 4\pi \displaystyle\int_0^{2ah+h^2} \sqrt{u}\left(\frac{1}{2}\, du\right) = 2\pi\left[\frac{2}{3}u^{3/2}\right]_0^{2ah+h^2} = \frac{4}{3}\pi\left(2ah + h^2\right)^{3/2}$.

12. A shell has radius x, circumference $2\pi x$, and height $\tan x - x$.



$V = \int_0^{\pi/3} 2\pi x\,(\tan x - x)\, dx$

13. A shell has radius $\frac{\pi}{2} - x$, circumference $2\pi\left(\frac{\pi}{2} - x\right)$, and height $\cos^2 x - \frac{1}{4}$.



$y = \cos^2 x$ intersects $y = \frac{1}{4}$ when $\cos^2 x = \frac{1}{4} \quad \Leftrightarrow$

$\cos x = \pm\frac{1}{2} \quad [\,|x| \le \pi/2] \quad \Leftrightarrow \quad x = \pm\frac{\pi}{3}$.

$$V = \int_{-\pi/3}^{\pi/3} 2\pi\left(\frac{\pi}{2} - x\right)\left(\cos^2 x - \frac{1}{4}\right) dx$$

14. A shell has radius $x - (-1) = x + 1$, circumference $2\pi(x + 1)$, and height $\ln x$.



$V = \int_1^4 2\pi(x + 1)\, \ln x\, dx$

15. $3x^2 = x^3 \quad \Leftrightarrow \quad x^3 - 3x^2 = 0 \quad \Leftrightarrow$

$x^2(x - 3) = 0 \quad \Leftrightarrow \quad x = 0$ or $x = 3$

$y = 3x^2 \quad \Rightarrow \quad x = \sqrt{\dfrac{y}{3}}\ (x > 0)$

$y = x^3 \quad \Rightarrow \quad x = \sqrt[3]{y}$



[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(a) With x as the variable of integration, we use the method of cylindrical shells.

$$V = \int_0^3 2\pi(x+1)(3x^2 - x^3)\,dx = \int_0^3 2\pi(2x^3 - x^4 + 3x^2)\,dx = 2\pi\left[\tfrac{1}{2}x^4 - \tfrac{1}{5}x^5 + x^3\right]_0^3$$
$$= 2\pi\left[\left(\frac{81}{2} - \frac{243}{5} + 27\right) - 0\right] = 2\pi \cdot \frac{189}{10} = \frac{189}{5}\pi$$

(b) With y as the variable of integration, we use washers with inner radius $\sqrt{\dfrac{y}{3}} + 1$ and outer radius $\sqrt[3]{y} + 1$.

The area of a cross-section is

$$\pi\left(\sqrt[3]{y} + 1\right)^2 - \pi\left(\sqrt{\frac{y}{3}} + 1\right)^2 = \pi\left[(y^{2/3} + 2y^{1/3} + 1) - \left(\frac{y}{3} + \frac{2}{\sqrt{3}}\,y^{1/2} + 1\right)\right]$$
$$= \pi\left(y^{2/3} + 2y^{1/3} - \frac{y}{3} - \frac{2}{\sqrt{3}}\,y^{1/2}\right)$$

$$V = \int_0^{27} \pi\left(y^{2/3} + 2y^{1/3} - \frac{y}{3} - \frac{2}{\sqrt{3}}\,y^{1/2}\right) dy = \pi\left[\frac{3}{5}y^{5/3} + \frac{3}{2}y^{4/3} - \frac{1}{6}y^2 - \frac{4}{3\sqrt{3}}\,y^{3/2}\right]_0^{27}$$
$$= \pi\left[\left(\frac{729}{5} + \frac{243}{2} - \frac{243}{2} - 108\right) - 0\right] = \frac{189}{5}\pi$$

16. $\sqrt{x} = x^2 \quad \Rightarrow \quad x = x^4 \quad \Leftrightarrow \quad x^4 - x = 0 \quad \Leftrightarrow$

$x(x^3 - 1) = 0 \quad \Leftrightarrow \quad x = 0$ or $x = 1$

$y = \sqrt{x} \quad \Rightarrow \quad x = y^2 \ (x > 0)$

$y = x^2 \ (x > 0) \quad \Rightarrow \quad x = \sqrt{y}$



(a) With x as the variable of integration, we use washers with inner radius $3 - \sqrt{x}$ and outer radius $3 - x^2$. The area of a cross-section is $\pi(3 - x^2)^2 - \pi\left(3 - \sqrt{x}\right)^2 = \pi[(9 - 6x^2 + x^4) - (9 - 6x^{1/2} + x)] = \pi(-6x^2 + x^4 + 6x^{1/2} - x)$.

$$V = \int_0^1 \pi(-6x^2 + x^4 + 6x^{1/2} - x)\,dx = \pi\left[-2x^3 + \tfrac{1}{5}x^5 + 4x^{3/2} - \tfrac{1}{2}x^2\right]_0^1 = \pi\left[\left(-2 + \tfrac{1}{5} + 4 - \tfrac{1}{2}\right) - 0\right] = \tfrac{17}{10}\pi$$

(b) With y as the variable of integration, we use the method of cylindrical shells.

$$V = \int_0^1 2\pi(3 - y)\left(\sqrt{y} - y^2\right) dy = \int_0^1 2\pi(3y^{1/2} - 3y^2 - y^{3/2} + y^3)\,dy$$
$$= 2\pi\left[2y^{3/2} - y^3 - \tfrac{2}{5}y^{5/2} + \tfrac{1}{4}y^4\right]_0^1 = 2\pi\left[\left(2 - 1 - \tfrac{2}{5} + \tfrac{1}{4}\right) - 0\right] = 2\pi \cdot \tfrac{17}{20} = \tfrac{17}{10}\pi$$

17. (a) A cross-section is a washer with inner radius x^2 and outer radius x.

$V = \int_0^1 \pi\left[(x)^2 - (x^2)^2\right] dx = \int_0^1 \pi(x^2 - x^4)\,dx = \pi\left[\tfrac{1}{3}x^3 - \tfrac{1}{5}x^5\right]_0^1 = \pi\left[\tfrac{1}{3} - \tfrac{1}{5}\right] = \tfrac{2}{15}\pi$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) A cross-section is a washer with inner radius y and outer radius $\sqrt{y}$.

$$V = \int_0^1 \pi\left[\left(\sqrt{y}\right)^2 - y^2\right] dy = \int_0^1 \pi(y - y^2)\, dy = \pi\left[\tfrac{1}{2}y^2 - \tfrac{1}{3}y^3\right]_0^1 = \pi\left[\tfrac{1}{2} - \tfrac{1}{3}\right] = \tfrac{\pi}{6}$$

(c) A cross-section is a washer with inner radius $2 - x$ and outer radius $2 - x^2$.

$$V = \int_0^1 \pi\left[(2 - x^2)^2 - (2 - x)^2\right] dx = \int_0^1 \pi(x^4 - 5x^2 + 4x)\, dx = \pi\left[\tfrac{1}{5}x^5 - \tfrac{5}{3}x^3 + 2x^2\right]_0^1 = \pi\left[\tfrac{1}{5} - \tfrac{5}{3} + 2\right] = \tfrac{8}{15}\pi$$

18. (a) $A = \int_0^1 (2x - x^2 - x^3)\, dx = \left[x^2 - \tfrac{1}{3}x^3 - \tfrac{1}{4}x^4\right]_0^1 = 1 - \tfrac{1}{3} - \tfrac{1}{4} = \tfrac{5}{12}$

(b) A cross-section is a washer with inner radius x^3 and outer radius $2x - x^2$, so its area is $\pi(2x - x^2)^2 - \pi(x^3)^2$.

$$\begin{aligned} V &= \int_0^1 A(x)\, dx = \int_0^1 \pi[(2x - x^2)^2 - (x^3)^2]\, dx = \int_0^1 \pi(4x^2 - 4x^3 + x^4 - x^6)\, dx \\ &= \pi\left[\tfrac{4}{3}x^3 - x^4 + \tfrac{1}{5}x^5 - \tfrac{1}{7}x^7\right]_0^1 = \pi\left(\tfrac{4}{3} - 1 + \tfrac{1}{5} - \tfrac{1}{7}\right) = \tfrac{41}{105}\pi \end{aligned}$$

(c) Using the method of cylindrical shells,

$$V = \int_0^1 2\pi x(2x - x^2 - x^3)\, dx = \int_0^1 2\pi(2x^2 - x^3 - x^4)\, dx = 2\pi\left[\tfrac{2}{3}x^3 - \tfrac{1}{4}x^4 - \tfrac{1}{5}x^5\right]_0^1 = 2\pi\left(\tfrac{2}{3} - \tfrac{1}{4} - \tfrac{1}{5}\right) = \tfrac{13}{30}\pi.$$

19. (a) Using the Midpoint Rule on $[0, 1]$ with $f(x) = \tan(x^2)$ and $n = 4$, we estimate

$$A = \int_0^1 \tan(x^2)\, dx \approx \tfrac{1}{4}\left[\tan\left(\left(\tfrac{1}{8}\right)^2\right) + \tan\left(\left(\tfrac{3}{8}\right)^2\right) + \tan\left(\left(\tfrac{5}{8}\right)^2\right) + \tan\left(\left(\tfrac{7}{8}\right)^2\right)\right] \approx \tfrac{1}{4}(1.53) \approx 0.38$$

(b) Using the Midpoint Rule on $[0, 1]$ with $f(x) = \pi\tan^2(x^2)$ (for disks) and $n = 4$, we estimate

$$V = \int_0^1 f(x)\, dx \approx \tfrac{1}{4}\pi\left[\tan^2\left(\left(\tfrac{1}{8}\right)^2\right) + \tan^2\left(\left(\tfrac{3}{8}\right)^2\right) + \tan^2\left(\left(\tfrac{5}{8}\right)^2\right) + \tan^2\left(\left(\tfrac{7}{8}\right)^2\right)\right] \approx \tfrac{\pi}{4}(1.114) \approx 0.87$$

20. (a)



From the graph, we see that the curves intersect at $x = 0$ and at $x = a \approx 0.75$, with $1 - x^2 > x^6 - x + 1$ on $(0, a)$.

(b) The area of $\mathcal{R}$ is $A = \int_0^a \left[(1 - x^2) - (x^6 - x + 1)\right] dx = \left[-\tfrac{1}{3}x^3 - \tfrac{1}{7}x^7 + \tfrac{1}{2}x^2\right]_0^a \approx 0.12$.

(c) Using washers, the volume generated when $\mathcal{R}$ is rotated about the x-axis is

$$\begin{aligned} V &= \pi\textstyle\int_0^a [(1 - x^2)^2 - (x^6 - x + 1)^2]\, dx = \pi\int_0^a (-x^{12} + 2x^7 - 2x^6 + x^4 - 3x^2 + 2x)\, dx \\ &= \pi\left[-\tfrac{1}{13}x^{13} + \tfrac{1}{4}x^8 - \tfrac{2}{7}x^7 + \tfrac{1}{5}x^5 - x^3 + x^2\right]_0^a \approx 0.54 \end{aligned}$$

(d) Using shells, the volume generated when $\mathcal{R}$ is rotated about the y-axis is

$$V = \int_0^a 2\pi x[(1 - x^2) - (x^6 - x + 1)]\, dx = 2\pi\int_0^a (-x^3 - x^7 + x^2)\, dx = 2\pi\left[-\tfrac{1}{4}x^4 - \tfrac{1}{8}x^8 + \tfrac{1}{3}x^3\right]_0^a \approx 0.31.$$

21. $\int_0^{\pi/2} 2\pi x\cos x\, dx = \int_0^{\pi/2} (2\pi x)\cos x\, dx$

The solid is obtained by rotating the region $\mathcal{R} = \left\{(x, y) \mid 0 \le x \le \tfrac{\pi}{2}, 0 \le y \le \cos x\right\}$ about the y-axis.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

22. $\int_0^{\pi/2} 2\pi\cos^2 x\,dx = \int_0^{\pi/2} \pi\big(\sqrt{2}\cos x\big)^2\,dx$

The solid is obtained by rotating the region $\mathcal{R} = \big\{(x,y) \mid 0 \le x \le \frac{\pi}{2}, 0 \le y \le \sqrt{2}\,\cos x\big\}$ about the x-axis.

23. $\int_0^{\pi} \pi(2-\sin x)^2\,dx$

The solid is obtained by rotating the region $\mathcal{R} = \{(x,y) \mid 0 \le x \le \pi, 0 \le y \le 2-\sin x\}$ about the x-axis.

24. $\int_0^4 2\pi(6-y)(4y-y^2)\,dy$

The solid is obtained by rotating the region $\mathcal{R} = \big\{(x,y) \mid 0 \le x \le 4y-y^2, 0 \le y \le 4\big\}$ about the line $y = 6$.

25. Take the base to be the disk $x^2+y^2 \le 9$. Then $V = \int_{-3}^{3} A(x)\,dx$, where $A(x_0)$ is the area of the isosceles right triangle whose hypotenuse lies along the line $x = x_0$ in the xy-plane. The length of the hypotenuse is $2\sqrt{9-x^2}$ and the length of each leg is $\sqrt{2}\sqrt{9-x^2}$. $A(x) = \frac{1}{2}\big(\sqrt{2}\sqrt{9-x^2}\big)^2 = 9-x^2$, so

$$V = 2\int_0^3 A(x)\,dx = 2\int_0^3 (9-x^2)\,dx = 2\big[9x - \tfrac{1}{3}x^3\big]_0^3 = 2(27-9) = 36$$

26. $V = \int_{-1}^{1} A(x)\,dx = 2\int_0^1 A(x)\,dx = 2\int_0^1 \big[(2-x^2)-x^2\big]^2\,dx = 2\int_0^1 \big[2(1-x^2)\big]^2\,dx$

$= 8\int_0^1 (1-2x^2+x^4)\,dx = 8\big[x - \frac{2}{3}x^3 + \frac{1}{5}x^5\big]_0^1 = 8\big(1-\frac{2}{3}+\frac{1}{5}\big) = \frac{64}{15}$

27. Equilateral triangles with sides measuring $\frac{1}{4}x$ meters have height $\frac{1}{4}x\sin 60^\circ = \frac{\sqrt{3}}{8}x$. Therefore,

$A(x) = \frac{1}{2}\cdot\frac{1}{4}x\cdot\frac{\sqrt{3}}{8}x = \frac{\sqrt{3}}{64}x^2$. $\quad V = \int_0^{20} A(x)\,dx = \frac{\sqrt{3}}{64}\int_0^{20} x^2\,dx = \frac{\sqrt{3}}{64}\big[\frac{1}{3}x^3\big]_0^{20} = \frac{8000\sqrt{3}}{64\cdot 3} = \frac{125\sqrt{3}}{3}\ \text{m}^3$.

28. (a) By the symmetry of the problem, we consider only the solid to the right of the origin. The semicircular cross-sections perpendicular to the x-axis have radius $1-x$, so $A(x) = \frac{1}{2}\pi(1-x)^2$. Now we can calculate

$V = 2\int_0^1 A(x)\,dx = 2\int_0^1 \frac{1}{2}\pi(1-x)^2\,dx = \int_0^1 \pi(1-x)^2\,dx = -\frac{\pi}{3}\big[(1-x)^3\big]_0^1 = \frac{\pi}{3}$.

(b) Cut the solid with a plane perpendicular to the x-axis and passing through the y-axis. Fold the half of the solid in the region $x \le 0$ under the xy-plane so that the point $(-1,0)$ comes around and touches the point $(1,0)$. The resulting solid is a right circular cone of radius 1 with vertex at $(x,y,z) = (1,0,0)$ and with its base in the yz-plane, centered at the origin. The volume of this cone is $\frac{1}{3}\pi r^2 h = \frac{1}{3}\pi\cdot 1^2\cdot 1 = \frac{\pi}{3}$.

29. $f(x) = kx \;\Rightarrow\; 30\ \text{N} = k(15-12)\ \text{cm} \;\Rightarrow\; k = 10\ \text{N/cm} = 1000\ \text{N/m}$. $20\ \text{cm} - 12\ \text{cm} = 0.08\ \text{m} \;\Rightarrow$

$W = \int_0^{0.08} kx\,dx = 1000\int_0^{0.08} x\,dx = 500\big[x^2\big]_0^{0.08} = 500(0.08)^2 = 3.2\ \text{N·m} = 3.2\ \text{J}$.

30. The work needed to raise the elevator alone is $1600\ \text{lb} \times 30\ \text{ft} = 48{,}000$ ft-lb. The work needed to raise the bottom 170 ft of cable is $170\ \text{ft} \times 10\ \text{lb/ft} \times 30\ \text{ft} = 51{,}000$ ft-lb. The work needed to raise the top 30 ft of cable is $\int_0^{30} 10x\,dx = \big[5x^2\big]_0^{30} = 5\cdot 900 = 4500$ ft-lb. Adding these, we see that the total work needed is $48{,}000 + 51{,}000 + 4{,}500 = 103{,}500$ ft-lb.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

31. (a) The parabola has equation $y = ax^2$ with vertex at the origin and passing through $(4, 4)$. $4 = a \cdot 4^2 \;\Rightarrow\; a = \frac{1}{4} \;\Rightarrow\; y = \frac{1}{4}x^2 \;\Rightarrow\; x^2 = 4y \;\Rightarrow\; x = 2\sqrt{y}$. Each circular disk has radius $2\sqrt{y}$ and is moved $4 - y$ ft.



$$W = \int_0^4 \pi \left(2\sqrt{y}\right)^2 62.5(4 - y)\, dy = 250\pi \int_0^4 y(4 - y)\, dy$$

$$= 250\pi\left[2y^2 - \tfrac{1}{3}y^3\right]_0^4 = 250\pi\left(32 - \tfrac{64}{3}\right) = \tfrac{8000\pi}{3} \approx 8378 \text{ ft-lb}$$

(b) In part (a) we knew the final water level (0) but not the amount of work done. Here we use the same equation, except with the work fixed, and the lower limit of integration (that is, the final water level — call it h) unknown: $W = 4000 \;\Leftrightarrow\; 250\pi\left[2y^2 - \frac{1}{3}y^3\right]_h^4 = 4000 \;\Leftrightarrow\; \frac{16}{\pi} = \left[\left(32 - \frac{64}{3}\right) - \left(2h^2 - \frac{1}{3}h^3\right)\right] \;\Leftrightarrow\; h^3 - 6h^2 + 32 - \frac{48}{\pi} = 0$. We graph the function $f(h) = h^3 - 6h^2 + 32 - \frac{48}{\pi}$ on the interval $[0, 4]$ to see where it is 0. From the graph, $f(h) = 0$ for $h \approx 2.1$. So the depth of water remaining is about 2.1 ft.



32. A horizontal slice of cooking oil Δx m thick has a volume of $\pi r^2 h = \pi \cdot 2^2 \cdot \Delta x \text{ m}^3$, a mass of $920(4\pi\,\Delta x)$ kg, weighs about $(9.8)(3680\pi\,\Delta x) = 36{,}064\pi\Delta x$ N, and thus requires about $36{,}064\pi x_i^*\Delta x$ J of work for its removal (where $3 \le x_i^* \le 6$). The total work needed to empty the tank is

$$W = \lim_{n\to\infty} \sum_{i=1}^{n} 36{,}064\pi x_i^*\,\Delta x = \int_3^6 36{,}064\pi x\, dx = 36{,}064\pi\left[\tfrac{1}{2}x^2\right]_3^6 = 18{,}032\pi(36 - 9) = 486{,}864\pi \approx 1.53 \times 10^6 \text{ J}.$$

33. $f_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b f(t)\, dt = \frac{1}{\pi/4 - 0}\int_0^{\pi/4} \sec^2 t\, dt = \frac{4}{\pi}\left[\tan t\right]_0^{\pi/4} = \frac{4}{\pi}(1 - 0) = \frac{4}{\pi}$

34. (a) $f_{\text{avg}} = \dfrac{1}{b-a}\displaystyle\int_a^b f(x)\, dx = \frac{1}{4-1}\int_1^4 \frac{1}{\sqrt{x}}\, dx$

$$= \frac{1}{3}\int_1^4 x^{-1/2}\, dx = \frac{1}{3}\left[2\sqrt{x}\right]_1^4$$

$$= \tfrac{2}{3}(2 - 1) = \tfrac{2}{3}$$

(b) $f(c) = f_{\text{avg}} \;\Leftrightarrow\; \dfrac{1}{\sqrt{c}} = \dfrac{2}{3} \;\Leftrightarrow\; \sqrt{c} = \dfrac{3}{2} \;\Leftrightarrow\; c = \dfrac{9}{4}$

(c)



35. (a) The regions $\mathcal{R}_1$ and $\mathcal{R}_2$ are shown in the figure.



The area of $\mathcal{R}_1$ is $A_1 = \int_0^b x^2\, dx = \left[\frac{1}{3}x^3\right]_0^b = \frac{1}{3}b^3$, and the area of $\mathcal{R}_2$ is $A_2 = \int_0^{b^2} \sqrt{y}\, dy = \left[\frac{2}{3}y^{3/2}\right]_0^{b^2} = \frac{2}{3}b^3$. So there is no solution to $A_1 = A_2$ for $b \neq 0$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(b) Using disks, we calculate the volume of rotation of $\mathcal{R}_1$ about the x-axis to be $V_{1,x} = \pi \int_0^b (x^2)^2\,dx = \frac{1}{5}\pi b^5$.

Using cylindrical shells, we calculate the volume of rotation of $\mathcal{R}_1$ about the y-axis to be

$V_{1,y} = 2\pi \int_0^b x(x^2)\,dx = 2\pi\left[\frac{1}{4}x^4\right]_0^b = \frac{1}{2}\pi b^4$. So $V_{1,x} = V_{1,y} \quad \Leftrightarrow \quad \frac{1}{5}\pi b^5 = \frac{1}{2}\pi b^4 \quad \Leftrightarrow \quad 2b = 5 \quad \Leftrightarrow \quad b = \frac{5}{2}$.

So the volumes of rotation about the x- and y-axes are the same for $b = \frac{5}{2}$.

(c) We use cylindrical shells to calculate the volume of rotation of $\mathcal{R}_2$ about the x-axis:

$\mathcal{R}_{2,x} = 2\pi \displaystyle\int_0^{b^2} y\left(\sqrt{y}\right)\,dy = 2\pi\left[\frac{2}{5}y^{5/2}\right]_0^{b^2} = \frac{4}{5}\pi b^5$. We already know the volume of rotation of $\mathcal{R}_1$ about the x-axis from part (b), and $\mathcal{R}_{1,x} = \mathcal{R}_{2,x} \quad \Leftrightarrow \quad \frac{1}{5}\pi b^5 = \frac{4}{5}\pi b^5$, which has no solution for $b \neq 0$.

(d) We use disks to calculate the volume of rotation of $\mathcal{R}_2$ about the y-axis: $\mathcal{R}_{2,y} = \pi \int_0^{b^2} \left(\sqrt{y}\right)^2\,dy = \pi\left[\frac{1}{2}y^2\right]_0^{b^2} = \frac{1}{2}\pi b^4$.

We know the volume of rotation of $\mathcal{R}_1$ about the y-axis from part (b), and $\mathcal{R}_{1,y} = \mathcal{R}_{2,y} \quad \Leftrightarrow \quad \frac{1}{2}\pi b^4 = \frac{1}{2}\pi b^4$. But this equation is true for all b, so the volumes of rotation about the y-axis are equal for all values of b.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

PROBLEMS PLUS

1. The volume generated from $x = 0$ to $x = b$ is $\int_0^b \pi[f(x)]^2\,dx$. Hence, we are given that $b^2 = \int_0^b \pi[f(x)]^2\,dx$ for all $b > 0$. Differentiating both sides of this equation with respect to b using the Fundamental Theorem of Calculus gives $2b = \pi[f(b)]^2 \quad \Rightarrow \quad f(b) = \sqrt{2b/\pi}$, since f is positive. Therefore, $f(x) = \sqrt{2x/\pi}$.

2. The total area of the region bounded by the parabola $y = x - x^2 = x(1 - x)$ and the x-axis is $\int_0^1 (x - x^2)\,dx = \left[\frac{1}{2}x^2 - \frac{1}{3}x^3\right]_0^1 = \frac{1}{6}$. Let the slope of the line we are looking for be m. Then the area above this line but below the parabola is $\int_0^a \left[(x - x^2) - mx\right]dx$, where a is the x-coordinate of the point of intersection of the line and the parabola. We find the point of intersection by solving the equation $x - x^2 = mx \quad \Leftrightarrow \quad 1 - x = m \quad \Leftrightarrow \quad x = 1 - m$. So the value of a is $1 - m$, and



$$\begin{aligned}\int_0^{1-m} \left[(x - x^2) - mx\right]dx &= \int_0^{1-m} \left[(1 - m)x - x^2\right]dx = \left[\tfrac{1}{2}(1 - m)x^2 - \tfrac{1}{3}x^3\right]_0^{1-m} \\ &= \tfrac{1}{2}(1 - m)(1 - m)^2 - \tfrac{1}{3}(1 - m)^3 = \tfrac{1}{6}(1 - m)^3\end{aligned}$$

We want this to be half of $\frac{1}{6}$, so $\frac{1}{6}(1 - m)^3 = \frac{1}{12} \quad \Leftrightarrow \quad (1 - m)^3 = \frac{6}{12} \quad \Leftrightarrow \quad 1 - m = \sqrt[3]{\frac{1}{2}} \quad \Leftrightarrow \quad m = 1 - \frac{1}{\sqrt[3]{2}}$. So the slope of the required line is $1 - \frac{1}{\sqrt[3]{2}} \approx 0.206$.

3. We must find expressions for the areas A and B, and then set them equal and see what this says about the curve C. If $P = \left(a, 2a^2\right)$, then area A is just $\int_0^a (2x^2 - x^2)\,dx = \int_0^a x^2\,dx = \frac{1}{3}a^3$. To find area B, we use y as the variable of integration. So we find the equation of the middle curve as a function of y: $y = 2x^2 \quad \Leftrightarrow \quad x = \sqrt{y/2}$, since we are concerned with the first quadrant only. We can express area B as

$$\int_0^{2a^2} \left[\sqrt{y/2} - C(y)\right]dy = \left[\frac{4}{3}(y/2)^{3/2}\right]_0^{2a^2} - \int_0^{2a^2} C(y)\,dy = \frac{4}{3}a^3 - \int_0^{2a^2} C(y)\,dy$$

where $C(y)$ is the function with graph C. Setting $A = B$, we get $\frac{1}{3}a^3 = \frac{4}{3}a^3 - \int_0^{2a^2} C(y)\,dy \quad \Leftrightarrow \quad \int_0^{2a^2} C(y)\,dy = a^3$. Now we differentiate this equation with respect to a using the Chain Rule and the Fundamental Theorem:

$C(2a^2)(4a) = 3a^2 \quad \Rightarrow \quad C(y) = \frac{3}{4}\sqrt{y/2}$, where $y = 2a^2$. Now we can solve for y: $x = \frac{3}{4}\sqrt{y/2} \quad \Rightarrow$

$x^2 = \frac{9}{16}(y/2) \quad \Rightarrow \quad y = \frac{32}{9}x^2$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

4. (a) Take slices perpendicular to the line through the center C of the bottom of the glass and the point P where the top surface of the water meets the bottom of the glass.



A typical rectangular cross-section y units above the axis of the glass has width $2\,|QR| = 2\sqrt{r^2-y^2}$ and length $h = |QS| = \dfrac{L}{2r}\,(r-y)$. [Triangles PQS and PAB are similar, so $\dfrac{h}{L} = \dfrac{|PQ|}{|PA|} = \dfrac{r-y}{2r}$.] Thus,

$$\begin{aligned} V &= \int_{-r}^{r} 2\sqrt{r^2-y^2}\cdot\frac{L}{2r}\,(r-y)\,dy = L\int_{-r}^{r}\left(1-\frac{y}{r}\right)\sqrt{r^2-y^2}\,dy \\ &= L\int_{-r}^{r}\sqrt{r^2-y^2}\,dy - \frac{L}{r}\int_{-r}^{r} y\sqrt{r^2-y^2}\,dy \\ &= L\cdot\frac{\pi r^2}{2} - \frac{L}{r}\cdot 0 \qquad \left[\begin{array}{c}\text{the first integral is the area of a semicircle of radius } r,\\ \text{and the second has an odd integrand}\end{array}\right] \qquad = \frac{\pi r^2 L}{2} \end{aligned}$$

(b) Slice parallel to the plane through the axis of the glass and the point of contact P. (This is the plane determined by P, B, and C in the figure.) $STUV$ is a typical trapezoidal slice. With respect to an x-axis with origin at C as shown, if S and V have x-coordinate x, then $|SV| = 2\sqrt{r^2-x^2}$. Projecting the trapezoid $STUV$ onto the plane of the triangle PAB (call the projection $S'T'U'V'$), we see that $|AP| = 2r$, $|SV| = 2\sqrt{r^2-x^2}$, and $|S'P| = |V'A| = \frac{1}{2}(|AP| - |SV|) = r - \sqrt{r^2-x^2}$.



By similar triangles, $\dfrac{|ST|}{|S'P|} = \dfrac{|AB|}{|AP|}$, so $|ST| = \left(r - \sqrt{r^2-x^2}\right)\cdot\dfrac{L}{2r}$. In the same way, we find that $\dfrac{|VU|}{|V'P|} = \dfrac{|AB|}{|AP|}$, so $|VU| = |V'P|\cdot\dfrac{L}{2r} = (|AP| - |V'A|)\cdot\dfrac{L}{2r} = \left(r + \sqrt{r^2-x^2}\right)\cdot\dfrac{L}{2r}$. The area $A(x)$ of the trapezoid $STUV$ is $\frac{1}{2}|SV|\cdot(|ST| + |VU|)$; that is,

$A(x) = \frac{1}{2}\cdot 2\sqrt{r^2-x^2}\cdot\left[\left(r-\sqrt{r^2-x^2}\right)\cdot\dfrac{L}{2r} + \left(r+\sqrt{r^2-x^2}\right)\cdot\dfrac{L}{2r}\right] = L\sqrt{r^2-x^2}$. Thus,

$$V = \int_{-r}^{r} A(x)\,dx = L\int_{-r}^{r}\sqrt{r^2-x^2}\,dx = L\cdot\frac{\pi r^2}{2} = \frac{\pi r^2 L}{2}.$$

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) See the computation of V in part (a) or part (b).

(d) The volume of the water is exactly half the volume of the cylindrical glass, so $V = \frac{1}{2}\pi r^2 L$.

(e) Choose x-, y-, and z-axes as shown in the figure. Then slices perpendicular to the x-axis are triangular, slices perpendicular to the y-axis are rectangular, and slices perpendicular to the z-axis are segments of circles. Using triangular slices, we find that the area $A(x)$ of a typical slice DEF, where D has x-coordinate x, is given by $A(x) = \frac{1}{2}|DE| \cdot |EF| = \frac{1}{2}|DE| \cdot \left(\frac{L}{r}\,|DE|\right) = \frac{L}{2r}\,|DE|^2 = \frac{L}{2r}\left(r^2 - x^2\right)$. Thus,





$$V = \int_{-r}^{r} A(x)\,dx = \frac{L}{2r}\int_{-r}^{r}(r^2 - x^2)\,dx = \frac{L}{r}\int_{-r}^{r}(r^2 - x^2)\,dx = \frac{L}{r}\left[r^2 x - \frac{x^3}{3}\right]_0^r$$

$$= \frac{L}{r}\left(r^3 - \frac{r^3}{3}\right) = \frac{L}{r}\cdot\tfrac{2}{3}r^3 = \tfrac{2}{3}r^2 L \qquad \text{[This is } 2/(3\pi) \approx 0.21 \text{ of the volume of the glass.]}$$

5. We are given that the rate of change of the volume of water is $\dfrac{dV}{dt} = -kA(x)$, where k is some positive constant and $A(x)$ is the area of the surface when the water has depth x. Now we are concerned with the rate of change of the depth of the water with respect to time, that is, $\dfrac{dx}{dt}$. But by the Chain Rule, $\dfrac{dV}{dt} = \dfrac{dV}{dx}\dfrac{dx}{dt}$, so the first equation can be written $\dfrac{dV}{dx}\dfrac{dx}{dt} = -kA(x)$ $(\star)$. Also, we know that the total volume of water up to a depth x is $V(x) = \int_0^x A(s)\,ds$, where $A(s)$ is the area of a cross-section of the water at a depth s. Differentiating this equation with respect to x, we get $dV/dx = A(x)$. Substituting this into equation $\star$, we get $A(x)(dx/dt) = -kA(x) \quad \Rightarrow \quad dx/dt = -k$, a constant.

6. (a) The volume above the surface is $\int_0^{L-h} A(y)\,dy = \int_{-h}^{L-h} A(y)\,dy - \int_{-h}^{0} A(y)\,dy$. So the proportion of volume above the surface is $\dfrac{\int_0^{L-h} A(y)\,dy}{\int_{-h}^{L-h} A(y)\,dy} = \dfrac{\int_{-h}^{L-h} A(y)\,dy - \int_{-h}^{0} A(y)\,dy}{\int_{-h}^{L-h} A(y)\,dy}$. Now by Archimedes' Principle, we have $F = W \quad \Rightarrow$ $\rho_f g \int_{-h}^{0} A(y)\,dy = \rho_0 g \int_{-h}^{L-h} A(y)\,dy$, so $\int_{-h}^{0} A(y)\,dy = \left(\rho_0/\rho_f\right)\int_{-h}^{L-h} A(y)\,dy$. Therefore, $\dfrac{\int_0^{L-h} A(y)\,dy}{\int_{-h}^{L-h} A(y)\,dy} = \dfrac{\int_{-h}^{L-h} A(y)\,dy - \left(\rho_0/\rho_f\right)\int_{-h}^{L-h} A(y)\,dy}{\int_{-h}^{L-h} A(y)\,dy} = \dfrac{\rho_f - \rho_0}{\rho_f}$, so the percentage of volume above the surface is $100\left(\dfrac{\rho_f - \rho_0}{\rho_f}\right)\%$.

(b) For an iceberg, the percentage of volume above the surface is $100\left(\frac{1030 - 917}{1030}\right)\% \approx 11\%$.

(c) No, the water does not overflow. Let V_i be the volume of the ice cube, and let V_w be the volume of the water which results from the melting. Then by the formula derived in part (a), the volume of ice above the surface of the water is

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$\big[(\rho_f-\rho_0)/\rho_f\big]V_i$, so the volume below the surface is $V_i-\big[(\rho_f-\rho_0)/\rho_f\big]V_i=(\rho_0/\rho_f)V_i$. Now the mass of the ice cube is the same as the mass of the water which is created when it melts, namely $m=\rho_0V_i=\rho_fV_w \;\Rightarrow$ $V_w=(\rho_0/\rho_f)V_i$. So when the ice cube melts, the volume of the resulting water is the same as the underwater volume of the ice cube, and so the water does not overflow.

(d) The figure shows the instant when the height of the exposed part of the ball is y.



Using the volume formula from Exercise 6.2.61, $V=\frac{1}{3}\pi h^2(3r-h)$, with $r=0.4$ and $h=0.8-y$, we see that the volume of the submerged part of the sphere is $\frac{1}{3}\pi(0.8-y)^2[1.2-(0.8-y)]$, so its weight is $1000g\cdot\frac{1}{3}\pi s^2(1.2-s)$, where $s=0.8-y$. Then the work done to submerge the sphere is

$$W=\int_0^{0.8} g\tfrac{1000}{3}\pi s^2(1.2-s)\,ds=g\tfrac{1000}{3}\pi\int_0^{0.8}(1.2s^2-s^3)\,ds$$
$$=g\tfrac{1000}{3}\pi\big[0.4s^3-\tfrac{1}{4}s^4\big]_0^{0.8}=g\tfrac{1000}{3}\pi(0.2048-0.1024)=9.8\,\tfrac{1000}{3}\,\pi(0.1024)\approx 1.05\times10^3\ \text{J}$$

7. A typical sphere of radius r is shown in the figure. We wish to maximize the shaded volume V, which can be thought of as the volume of a hemisphere of radius r minus the volume of the spherical cap with height $h=1-\sqrt{1-r^2}$ and radius 1.



$$V=\tfrac{1}{2}\cdot\tfrac{4}{3}\pi r^3-\tfrac{1}{3}\pi\big(1-\sqrt{1-r^2}\big)^2\big[3(1)-\big(1-\sqrt{1-r^2}\big)\big]\quad\text{[by Exercise 6.2.61]}$$
$$=\tfrac{1}{3}\pi\big[2r^3-\big(2-2\sqrt{1-r^2}-r^2\big)\big(2+\sqrt{1-r^2}\big)\big]$$
$$=\tfrac{1}{3}\pi\big[2r^3-2+\big(r^2+2\big)\sqrt{1-r^2}\big]$$

$$V'=\tfrac{1}{3}\pi\left[6r^2+\frac{\big(r^2+2\big)(-r)}{\sqrt{1-r^2}}+\sqrt{1-r^2}(2r)\right]=\tfrac{1}{3}\pi\left[\frac{6r^2\sqrt{1-r^2}-r\big(r^2+2\big)+2r\big(1-r^2\big)}{\sqrt{1-r^2}}\right]$$
$$=\tfrac{1}{3}\pi\left(\frac{6r^2\sqrt{1-r^2}-3r^3}{\sqrt{1-r^2}}\right)=\frac{\pi r^2\big(2\sqrt{1-r^2}-r\big)}{\sqrt{1-r^2}}$$

$V'(r)=0 \;\Leftrightarrow\; 2\sqrt{1-r^2}=r \;\Leftrightarrow\; 4-4r^2=r^2 \;\Leftrightarrow\; r^2=\frac{4}{5} \;\Leftrightarrow\; r=\frac{2}{\sqrt{5}}\approx0.89.$

Since $V'(r)>0$ for $0<r<\frac{2}{\sqrt{5}}$ and $V'(r)<0$ for $\frac{2}{\sqrt{5}}<r<1$, we know that V attains a maximum at $r=\frac{2}{\sqrt{5}}$.

8. We want to find the volume of that part of the sphere which is below the surface of the water. As we can see from the diagram, this region is a cap of a sphere with radius r and height $r+d$. If we can find an expression for d in terms of h, r and θ, then we can determine the volume of the region [see Exercise 6.2.61], and then differentiate with respect to r to find the maximum. We see that



$$\sin\theta=\frac{r}{h-d} \;\Leftrightarrow\; h-d=\frac{r}{\sin\theta} \;\Leftrightarrow\; d=h-r\csc\theta.$$

[continued]

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

Now we can use the formula from Exercise 6.2.61 to find the volume of water displaced:

$$V = \tfrac{1}{3}\pi h^2(3r-h) = \tfrac{1}{3}\pi(r+d)^2\,[3r-(r+d)] = \tfrac{1}{3}\pi(r+h-r\csc\theta)^2(2r-h+r\csc\theta)$$
$$= \tfrac{\pi}{3}[r(1-\csc\theta)+h]^2[r(2+\csc\theta)-h]$$

Now we differentiate with respect to r:

$$dV/dr = \tfrac{\pi}{3}\big([r(1-\csc\theta)+h]^2(2+\csc\theta)+2[r(1-\csc\theta)+h](1-\csc\theta)[r(2+\csc\theta)-h]\big)$$
$$= \tfrac{\pi}{3}[r(1-\csc\theta)+h]([r(1-\csc\theta)+h](2+\csc\theta)+2(1-\csc\theta)[r(2+\csc\theta)-h])$$
$$= \tfrac{\pi}{3}[r(1-\csc\theta)+h](3(2+\csc\theta)(1-\csc\theta)r+[(2+\csc\theta)-2(1-\csc\theta)]h)$$
$$= \tfrac{\pi}{3}[r(1-\csc\theta)+h][3(2+\csc\theta)(1-\csc\theta)r+3h\csc\theta]$$

This is 0 when $r = \dfrac{h}{\csc\theta-1}$ and when $r = \dfrac{h\csc\theta}{(\csc\theta+2)(\csc\theta-1)}$. Now since $V\left(\dfrac{h}{\csc\theta-1}\right)=0$ (the first factor vanishes; this corresponds to $d=-r$), the maximum volume of water is displaced when $r = \dfrac{h\csc\theta}{(\csc\theta-1)(\csc\theta+2)}$. (Our intuition tells us that a maximum value does exist, and it must occur at a critical number.) Multiplying numerator and denominator by $\sin^2\theta$, we get an alternative form of the answer: $r = \dfrac{h\sin\theta}{\sin\theta+\cos 2\theta}$.

9. (a) Stacking disks along the y-axis gives us $V = \int_0^h \pi\,[f(y)]^2\,dy$.

(b) Using the Chain Rule, $\dfrac{dV}{dt} = \dfrac{dV}{dh}\cdot\dfrac{dh}{dt} = \pi\,[f(h)]^2\,\dfrac{dh}{dt}$.

(c) $kA\sqrt{h} = \pi[f(h)]^2\,\dfrac{dh}{dt}$. Set $\dfrac{dh}{dt}=C$: $\pi[f(h)]^2\,C = kA\sqrt{h} \;\Rightarrow\; [f(h)]^2 = \dfrac{kA}{\pi C}\sqrt{h} \;\Rightarrow\; f(h) = \sqrt{\dfrac{kA}{\pi C}}\,h^{1/4}$; that is, $f(y) = \sqrt{\dfrac{kA}{\pi C}}\,y^{1/4}$. The advantage of having $\dfrac{dh}{dt}=C$ is that the markings on the container are equally spaced.

10. (a) We first use the cylindrical shell method to express the volume V in terms of h, r, and ω:

$$V = \int_0^r 2\pi xy\,dx = \int_0^r 2\pi x\left[h+\frac{\omega^2x^2}{2g}\right]dx = 2\pi\int_0^r\left(hx+\frac{\omega^2x^3}{2g}\right)dx$$
$$= 2\pi\left[\frac{hx^2}{2}+\frac{\omega^2x^4}{8g}\right]_0^r = 2\pi\left[\frac{hr^2}{2}+\frac{\omega^2r^4}{8g}\right] = \pi hr^2+\frac{\pi\omega^2r^4}{4g} \quad\Rightarrow$$

$h = \dfrac{V-\left(\pi\omega^2r^4\right)/(4g)}{\pi r^2} = \dfrac{4gV-\pi\omega^2r^4}{4\pi gr^2}$.

(b) The surface touches the bottom when $h=0 \;\Rightarrow\; 4gV-\pi\omega^2r^4=0 \;\Rightarrow\; \omega^2 = \dfrac{4gV}{\pi r^4} \;\Rightarrow\; \omega = \dfrac{2\sqrt{gV}}{\sqrt{\pi}r^2}$.

To spill over the top, $y(r) > L \;\Leftrightarrow$

$$L < h+\frac{\omega^2r^2}{2g} = \frac{4gV-\pi\omega^2r^4}{4\pi gr^2}+\frac{\omega^2r^2}{2g} = \frac{4gV}{4\pi gr^2}-\frac{\pi\omega^2r^2}{4\pi gr^2}+\frac{\omega^2r^2}{2g}$$
$$= \frac{V}{\pi r^2}-\frac{\omega^2r^2}{4g}+\frac{\omega^2r^2}{2g} = \frac{V}{\pi r^2}+\frac{\omega^2r^2}{4g} \quad\Leftrightarrow$$

$\dfrac{\omega^2r^2}{4g} > L-\dfrac{V}{\pi r^2} = \dfrac{\pi r^2L-V}{\pi r^2} \;\Leftrightarrow\; \omega^2 > \dfrac{4g\left(\pi r^2L-V\right)}{\pi r^4}$. So for spillage, the angular speed should be $\omega > \dfrac{2\sqrt{g(\pi r^2L-V)}}{r^2\sqrt{\pi}}$.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

(c) (i) Here we have $r = 2$, $L = 7$, $h = 7 - 5 = 2$. When $x = 1$, $y = 7 - 4 = 3$. Therefore, $3 = 2 + \dfrac{\omega^2 \cdot 1^2}{2 \cdot 32} \quad \Rightarrow$

$1 = \dfrac{\omega^2}{2 \cdot 32} \quad \Rightarrow \quad \omega^2 = 64 \quad \Rightarrow \quad \omega = 8$ rad/s. $V = \pi(2)(2)^2 + \dfrac{\pi \cdot 8^2 \cdot 2^4}{4g} = 8\pi + 8\pi = 16\pi$ ft^2.

(ii) At the wall, $x = 2$, so $y = 2 + \dfrac{8^2 \cdot 2^2}{2 \cdot 32} = 6$ and the surface is $7 - 6 = 1$ ft below the top of the tank.

11. The cubic polynomial passes through the origin, so let its equation be $y = px^3 + qx^2 + rx$. The curves intersect when $px^3 + qx^2 + rx = x^2 \quad \Leftrightarrow$ $px^3 + (q-1)x^2 + rx = 0$. Call the left side $f(x)$. Since $f(a) = f(b) = 0$, another form of f is



$$f(x) = px(x-a)(x-b) = px[x^2 - (a+b)x + ab]$$
$$= p[x^3 - (a+b)x^2 + abx]$$

Since the two areas are equal, we must have $\int_0^a f(x)\,dx = -\int_a^b f(x)\,dx \quad \Rightarrow$ $[F(x)]_0^a = [F(x)]_b^a \quad \Rightarrow \quad F(a) - F(0) = F(a) - F(b) \quad \Rightarrow \quad F(0) = F(b)$, where F is an antiderivative of f. Now $F(x) = \int f(x)\,dx = \int p[x^3 - (a+b)x^2 + abx]\,dx = p\left[\frac{1}{4}x^4 - \frac{1}{3}(a+b)x^3 + \frac{1}{2}abx^2\right] + C$, so $F(0) = F(b) \quad \Rightarrow \quad C = p\left[\frac{1}{4}b^4 - \frac{1}{3}(a+b)b^3 + \frac{1}{2}ab^3\right] + C \quad \Rightarrow \quad 0 = p\left[\frac{1}{4}b^4 - \frac{1}{3}(a+b)b^3 + \frac{1}{2}ab^3\right] \quad \Rightarrow$ $0 = 3b - 4(a+b) + 6a$ [multiply by $12/(pb^3)$, $b \neq 0$] $\Rightarrow \quad 0 = 3b - 4a - 4b + 6a \quad \Rightarrow \quad b = 2a$. Hence, b is twice the value of a.

12. (a) Place the round flat tortilla on an xy-coordinate system as shown in the first figure. An equation of the circle is $x^2 + y^2 = 4^2$ and the height of a cross-section is $2\sqrt{16 - x^2}$.



Now look at a cross-section with central angle θ_x as shown in the second figure (r is the radius of the circular cylinder). The filled area $A(x)$ is equal to the area $A_1(x)$ of the sector minus the area $A_2(x)$ of the triangle.



$$A(x) = A_1(x) - A_2(x) = \tfrac{1}{2}r^2\theta_x - \tfrac{1}{2}r^2 \sin\theta_x \qquad \text{[area formulas from trigonometry]}$$
$$= \tfrac{1}{2}r(r\theta_x) - \tfrac{1}{2}r^2 \sin\left(\frac{s}{r}\right) \qquad \text{[arc length } s = r\theta_x \quad \Rightarrow \quad \theta_x = s/r\text{]}$$
$$= \tfrac{1}{2}r \cdot 2\sqrt{16 - x^2} - \tfrac{1}{2}r^2 \sin\left(\frac{2\sqrt{16 - x^2}}{r}\right) \qquad [s = 2\sqrt{16 - x^2}\,]$$
$$= r\sqrt{16 - x^2} - \tfrac{1}{2}r^2 \sin\left(\frac{2}{r}\sqrt{16 - x^2}\right) \quad (\star)$$

Note that the central angle θ_x will be small near the ends of the tortilla; that is, when $|x| \approx 4$. But near the center of

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

the tortilla (when $|x| \approx 0$), the central angle θ_x may exceed $180°$. Thus, the sine of θ_x will be negative and the second term in $(\star)$ will be positive (actually adding area to the area of the sector). The volume of the taco can be found by integrating the cross-sectional areas from $x = -4$ to $x = 4$. Thus,

$$V(x) = \int_{-4}^{4} A(x)\,dx = \int_{-4}^{4} \left[r\sqrt{16 - x^2} - \tfrac{1}{2}r^2 \sin\left(\frac{2}{r}\sqrt{16 - x^2}\right)\right] dx$$

(b) To find the value of r that maximizes the volume of the taco, we can define the function

$$V(r) = \int_{-4}^{4} \left[r\sqrt{16 - x^2} - \tfrac{1}{2}r^2 \sin\left(\frac{2}{r}\sqrt{16 - x^2}\right)\right] dx$$

The figure shows a graph of $y = V(r)$ and $y = V'(r)$. The maximum volume of about 52.94 occurs when $r \approx 2.2912$.



13. We assume that P lies in the region of positive x. Since $y = x^3$ is an odd function, this assumption will not affect the result of the calculation. Let $P = (a, a^3)$. The slope of the tangent to the curve $y = x^3$ at P is $3a^2$, and so the equation of the tangent is $y - a^3 = 3a^2(x - a) \quad \Leftrightarrow \quad y = 3a^2x - 2a^3$.



We solve this simultaneously with $y = x^3$ to find the other point of intersection: $x^3 = 3a^2x - 2a^3 \quad \Leftrightarrow$ $(x - a)^2(x + 2a) = 0$. So $Q = (-2a, -8a^3)$ is the other point of intersection. The equation of the tangent at Q is $y - (-8a^3) = 12a^2[x - (-2a)] \quad \Leftrightarrow \quad y = 12a^2x + 16a^3$. By symmetry, this tangent will intersect the curve again at $x = -2(-2a) = 4a$. The curve lies above the first tangent, and

below the second, so we are looking for a relationship between $A = \int_{-2a}^{a} \left[x^3 - (3a^2x - 2a^3)\right] dx$ and

$B = \int_{-2a}^{4a} \left[(12a^2x + 16a^3) - x^3\right] dx$. We calculate $A = \left[\frac{1}{4}x^4 - \frac{3}{2}a^2x^2 + 2a^3x\right]_{-2a}^{a} = \frac{3}{4}a^4 - (-6a^4) = \frac{27}{4}a^4$, and

$B = \left[6a^2x^2 + 16a^3x - \frac{1}{4}x^4\right]_{-2a}^{4a} = 96a^4 - (-12a^4) = 108a^4$. We see that $B = 16A = 2^4A$. This is because our calculation of area B was essentially the same as that of area A, with a replaced by $-2a$, so if we replace a with $-2a$ in our expression for A, we get $\frac{27}{4}(-2a)^4 = 108a^4 = B$.

14. From the solution to Problem 11 in Problems Plus following Chapter 4, an equation of the normal line through P is $y - a^2 = -\dfrac{1}{2a}(x - a) \quad \Rightarrow \quad y = -\dfrac{1}{2a}x + \dfrac{1}{2} + a^2$, and the x-coordinate of Q is $x = -a - \dfrac{1}{2a}$. The area of $\mathcal{R}$ is given by

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.

$$A(a)=\int_{-a-1/(2a)}^{a}\left[\left(-\frac{1}{2a}x+\frac{1}{2}+a^2\right)-x^2\right]dx=\left[-\frac{1}{4a}x^2+\frac{1}{2}x+a^2x-\frac{1}{3}x^3\right]_{-a-1/(2a)}^{a}$$

$$=\left(-\frac{1}{4}a+\frac{1}{2}a+a^3-\frac{1}{3}a^3\right)-\left[-\frac{1}{4a}\left(-a-\frac{1}{2a}\right)^2+\frac{1}{2}\left(-a-\frac{1}{2a}\right)+a^2\left(-a-\frac{1}{2a}\right)-\frac{1}{3}\left(-a-\frac{1}{2a}\right)^3\right]$$

$$=\left(\frac{1}{4}a+\frac{2}{3}a^3\right)-\left[-\frac{1}{4a}\left(a^2+1+\frac{1}{4a^2}\right)-\frac{1}{2}a-\frac{1}{4a}-a^3-\frac{1}{2}a-\frac{1}{3}\left(-a^3-\frac{3}{2}a-\frac{3}{4a}-\frac{1}{8a^3}\right)\right]$$

$$=\left(\frac{1}{4}a+\frac{2}{3}a^3\right)-\left(-\frac{2}{3}a^3-\frac{1}{48a^3}-\frac{3}{4}a-\frac{1}{4a}\right)$$

$$=\frac{4}{3}a^3+\frac{1}{48a^3}+a+\frac{1}{4a}=\frac{64a^6+48a^4+12a^2+1}{48a^3}=\frac{(4a^2+1)^3}{48a^3}$$

$$A'(a)=\frac{48a^3\cdot 3(4a^2+1)^2\cdot 8a-(4a^2+1)^3\cdot 144a^2}{(48a^3)^2}=\frac{48\cdot 3a^2(4a^2+1)^2[a\cdot 8a-(4a^2+1)]}{48\cdot 48\cdot a^3\cdot a^3}$$

$$=\frac{(4a^2+1)^2(4a^2-1)}{16a^4}=0 \quad\Rightarrow\quad a^2=\frac{1}{4} \quad\Rightarrow\quad a=\frac{1}{2}\ (a>0)$$

Since $A'(a)<0$ for $0<a<\frac{1}{2}$ and $A'(a)>0$ for $a>\frac{1}{2}$, there is an absolute minimum when $a=\frac{1}{2}$ by the First Derivative Test for Absolute Extreme Values.

© 2021 Cengage Learning. All Rights Reserved. May not be scanned, copied, or duplicated, or posted to a publicly accessible website, in whole or in part.